RNA Interference Pathways in Fungi: Mechanisms and Functions

Shwu-Shin Chang, Zhenyu Zhang, and Yi Liu

Department of Physiology, The University of Texas Southwestern Medical Center, Dallas, Texas 75390; email: Yi.Liu@UTsouthwestern.edu

Annu. Rev. Microbiol. 2012. 66:305-23

First published online as a Review in Advance on June 28, 2012

The Annual Review of Microbiology is online at micro.annualreviews.org

This article's doi: 10.1146/annurev-micro-092611-150138

Copyright © 2012 by Annual Reviews. All rights reserved

0066-4227/12/1013-0305\$20.00

Keywords

small noncoding RNA, Schizosaccharomyces pombe, Neurospora

Abstract

RNA interference (RNAi) is a conserved eukaryotic gene regulatory mechanism that uses small noncoding RNAs to mediate posttranscriptional/ transcriptional gene silencing. The fission yeast *Schizosaccharomyces pombe* and the filamentous fungus *Neurospora crassa* have served as important model systems for RNAi research. Studies on these two organisms and other fungi have contributed significantly to our understanding of the mechanisms and functions of RNAi in eukaryotes. In addition, surprisingly diverse RNAimediated processes and small RNA biogenesis pathways have been discovered in fungi. In this review, we give an overview of different fungal RNAi pathways with a focus on their mechanisms and functions.

Contents

INTRODUCTION	306
QUELLING IN NEUROSPORA	307
The Discovery of Quelling	307
Molecular Mechanism of Quelling	307
qiRNA: A Type of sRNA Induced by DNA Damage	310
MEIOTIC SILENCING BY UNPAIRED DNA	310
RNAi-MEDIATED HETEROCHROMATIN FORMATION IN	
SCHIZOSACCHAROMYCES POMBE	312
RNAi IN BUDDING YEASTS	315
miRNA-LIKE sRNA IN NEUROSPORA	315
RNAi AS A HOST DEFENSE MECHANISM IN FUNGI	316
Transposon Control in Neurospora	316
An Antiviral Mechanism in Cryphonectria parasitica and Aspergillus nidulans	316
dsRNA/Viral Infection–Induced Transcriptional Response	
in Neurospora and C. parasitica	317
RNA SILENCING AND SMALL RNA STUDIES ON OTHER FILAMENTOUS	
FUNGI	317
Gene Silencing and Small RNAs in Mucor circinelloides	317
RNAi Studies on the Human Fungal Pathogen Cryptococcus neoformans	318
Small RNAs from <i>Magnaporthe oryzae</i>	318

INTRODUCTION

RNA interference (RNAi) in *Caenorhabditis elegans* was first described in 1998 as a phenomenon triggered by double-stranded RNA (dsRNA) that results in the silencing of the genes complementary to the dsRNA (31, 72). This process was initially thought to be a host defense mechanism against invading viruses and transposons (80); however, studies over the past decade have demonstrated that RNAi pathways use small noncoding RNAs (sRNAs) to regulate diverse cellular, developmental, and physiological processes. RNAi pathways are conserved in eukaryotes, and among the identified sRNAs, there are three major classes: small interfering RNA (siRNA), microRNA (miRNA), and PIWI-interacting RNA (piRNA) (10, 34, 54, 73, 100). Each class of sRNAs is produced differently and has diverse functions. In the siRNA and miRNA pathways, the RNase-III-like Dicer enzymes cleave dsRNA/hairpin RNA to generate 21- to 25-nt mature siRNA or miRNA duplexes. The duplex is incorporated into an effector complex, the RNA-induced silencing complex (RISC), containing an Argonaute protein with slicer activity. The passenger strand of the sRNA duplex is removed and the remaining guide RNA directs the RISC to complementary mRNA sequences, resulting in silencing by either mRNA cleavage or translational repression.

The fungi kingdom is a large and diverse group of eukaryotic organisms estimated to contain more than three million species, and has diverse and vital roles in the ecosystem (81). The RNAi-related phenomena were first described in plants and fungi (76, 85). Quelling, the RNAi-related posttranscriptional gene-silencing phenomenon in *Neurospora* was reported in 1992 (85). Since then, the identification of quelling components and the analysis of the pathway have contributed significantly to our understanding of RNAi-mediated gene silencing at the posttranscriptional level. RNAi was first found to be required for heterochromatin formation in the fission yeast

dsRNA:

double-stranded RNA

RNA interference

(RNAi): a conserved eukaryotic gene regulatory mechanism that uses small RNA to mediate gene silencing

sRNA: small noncoding RNA

microRNAs

(miRNAs): a type of small RNAs that are generated from precursor RNAs with stem-loop structures. microRNAs can silence gene expression at posttranscriptional and translational levels *Schizosaccharomyces pombe* (105), and this system remains the best-understood pathway of RNAimediated transcriptional gene silencing. In addition, RNAi components and their functions have been characterized in many fungal species (7). Furthermore, many classes of sRNAs have been found in different fungal species, revealing diverse sRNA biogenesis pathways. In this review, we focus on the mechanisms and functions of different fungal RNAi pathways.

QUELLING IN NEUROSPORA

The Discovery of Quelling

Soon after the discovery of cosuppression in plants (76), Romano & Macino (85) discovered a similar transgene-induced gene-silencing phenomenon in *Neurospora crassa* called quelling. Quelling was discovered by transforming *Neurospora* with DNA containing *albino-1* or *albino-3* genes, which are involved in carotenoid biosynthesis. After transformation, white/pale-yellow transformants with reduced *al* mRNA levels were obtained, indicating the silencing of endogenous *al* genes by the transgenes (9, 20, 32, 85). High copy numbers of the transgenes correlated with the quelling efficiency, suggesting that the presence of repetitive transgenes causes silencing. In addition, the quelled transformants spontaneously and progressively reverted to wild type or intermediate phenotypes owing to reductions in copy number of the transgene.

The phenotype of *al* quelled strains is dominant, suggesting that diffusible and *trans*-acting silencing molecules are involved in quelling (20). In addition, a sense RNA derived from the transgene was specifically detected in quelled strains, implying that aberrant transcription of transgenes is involved in silencing (20). Furthermore, quelling did not affect the levels of mRNA precursors, indicating that quelling is a posttranscriptional gene silencing (PTGS) mechanism (20). Together, these results led to the hypothesis that the production of aberrant RNA (aRNA) in the presence of multicopy transgenes causes PTGS.

Molecular Mechanism of Quelling

Cogoni & Macino (21) isolated *quelling-defective (qde)* mutants, which were mapped to three complementation groups, *qde-1*, *qde-2*, and *qde-3*. The cloning and studies of these *qde* genes contributed significantly to our understanding of an RNAi pathway that is well conserved across eukaryotes (13, 14, 21, 25). QDE-1, which encodes a cellular RNA-dependent RNA polymerase (RdRP), was the first eukaryotic RNAi component identified (22); it was discovered soon after the silencing effect of dsRNA was demonstrated in *C. elegans* (31). The requirement of an RdRP in quelling shows that dsRNA is an essential intermediate of PTGS in vivo.

QDE-2 encodes an Argonaute protein that is homologous to RDE-1 (RNAi-Deficient-1) in *C. elegans*, which is required for dsRNA-mediated silencing (11). The identification of QDE-2 provided the first evidence that RNAi in *C. elegans* and quelling share a common genetic mechanism.

QDE-3 belongs to the RecQ helicase family and is homologous to the mammalian Bloom's and Werner's syndrome proteins (23). RecQ helicases are involved in homologous recombination, DNA replication, and DNA repair. As expected, QDE-3 contributes to the DNA repair process in *Neurospora* (23, 51). The requirement of QDE-3 in quelling raises the possibility that repetitive transgenes may form aberrant DNA structures that are recognized by QDE-3 to promote aRNA and sRNA production. Interestingly, OsRecQ1, a RecQ helicase homolog in rice, is required for inverted-repeat-induced RNA silencing (16). rRecQ-1, a homolog of QDE-3 in rats, is associated with the piRNA-binding complex (58).

Expression of an inverted-repeat-containing transgene can bypass QDE-1 and QDE-3 to trigger gene silencing (14, 61), indicating that QDE-1 and QDE-3 function in producing aRNA

Quelling:

an RNAi-related gene-silencing mechanism in *Neurospora* that is induced by repetitive transgenic sequences. Quelling occurs in the vegetative growth stage

Heterochromatin:

chromosome regions that are tightly packed throughout the cell cycle and are mostly genetically inactive

PTGS:

posttranscriptional gene silencing

QDE-2:

Quelling-Deficient 2

and dsRNA from repetitive transgenes. In addition, these results suggest that QDE-1 is not involved in sRNA amplification. Although QDE-1 is expected to function as an RdRP to convert aRNA to dsRNA, it was not clear how aRNAs are produced and specifically recognized by QDE-1. In plants, RNA polymerases Pol IV and Pol V are required for production of the single-stranded noncoding RNA precursors of dsRNAs and siRNAs (71, 108). In the fission yeast, RNA Pol II produces the centromeric siRNA precursors (88). The structure of QDE-1 revealed that its catalytic core is similar to that of eukaryotic DNA-dependent RNA polymerase (DdRP) (87). In addition, QDE-1 is required for the production of ribosomal DNA (rDNA)-specific aRNA (61). Recently, we demonstrated that QDE-1 is both a DdRP and an RdRP and can use either single-stranded DNA or RNA (ssDNA or ssRNA) as templates to produce DNA/RNA hybrids and dsRNA, respectively (60, 61). Together, these results suggest that QDE-1 is an RNA polymerase that produces aRNA from ssDNA in the quelling pathway (61).

How is QDE-1 recruited to ssDNA substrates to produce aRNA? QDE-1 associates with RPA-1, which is a subunit of replication protein A (RPA), a ssDNA-binding complex involved in DNA replication, recombination, and repair (60, 78). Deletion of *rpa-3* in *Neurospora* abolished quelling and DNA-damage-induced production of QDE-2-interacting sRNAs (qiRNAs) (see below) (60). In addition, the interaction between QDE-1 and RPA requires QDE-3. QDE-1 can use ssDNA templates to produce dsRNA, an activity that is strongly promoted by RPA as it prevents the formation of DNA/RNA hybrids. These results suggest that RPA has two roles in the production of aRNA: It recruits QDE-1 to ssDNA with the aid of QDE-3, and it blocks the formation of DNA/RNA hybrids. This mechanism explains how aRNAs are produced and are specifically recognized by an RdRP in the quelling pathway. Repeat-induced gene silencing has been reported in many systems, including fungi, plants, *C. elegans*, *Drosophila melanogaster*, and mammals (46). The mechanism by which repetitive DNA sequences trigger the production of sRNA remains largely unclear. The involvement of QDE-3 and RPA in quelling suggests that DNA repeats may result in some aberrant DNA structures during DNA replication or recombination that may then be recognized by RNA polymerases to produce aRNAs and dsRNA.

Two partially redundant Neurospora Dicer-like proteins (DCL-1 and DCL-2) process dsRNA into ~25-nt sRNAs in an ATP-dependent manner (14). The deletion of both dicer genes completely abolishes quelling and Dicer activity. QDE-2 associates with siRNA duplexes and forms an inactive RISC complex (11, 12). In order to activate the RISC complex, the passenger strand of the siRNA duplex must be removed. We showed that QDE-2 and its slicer activity are required for single-stranded siRNA production, indicating that QDE-2 slices the passenger strand of siRNA to produce a nicked siRNA duplex (70); this reaction is analogous to the slicing of an mRNA substrate. This provides the first in vivo evidence that the Argonaute protein is involved in generating single-stranded siRNA. However, QDE-2 alone is not sufficient to remove the passenger strand. Biochemical purification of QDE-2 led to the identification of a QDE-2-interacting protein called QIP; this putative exonuclease is required for the removal of the passenger strand of siRNAs (70). Together, these results indicate that both the cleavage and removal of the passenger strand from the siRNA duplex are essential steps in RISC activation. A Drosophila ribonuclease, C3PO (component 3 promoter of RISC), was later found to promote RISC activation by removing siRNA passenger strand cleavage products (68), indicating that the RISC activation mechanism is similar in Neurospora and animals.

The current model for the *Neurospora* quelling pathway is depicted in **Figure 1**. Repetitive DNA sequences form aberrant DNA structures after encountering DNA replication stress. These aberrant DNA structures are recognized by QDE-3 and RPA, which recruit QDE-1 to the ssDNA locus. QDE-1 acts as a DdRP to produce aRNAs, which are then converted to dsRNAs by the RdRP activity of QDE-1. The long dsRNA precursors are processed into siRNA duplexes by

Annu. Rev. Microbiol. 2012.66:305-323. Downloaded from www.annualreviews.org by Universidade Federal do Parana on 03/05/13. For personal use only.





Figure 1

Models for RNAi-related pathways in *Neurospora crassa*. During the vegetative stage (*right*), both repetitive transgene (quelling) and ribosomal DNA loci (qiRNA) lead to the production of aRNAs by QDE-1 and QDE-3. Single-stranded aRNAs are then converted to dsRNA precursors by QDE-1. dsRNA are processed by DCLs into sRNAs, which are then loaded onto QDE-2/QIP-based RISC to execute gene silencing. Yellow arrows indicate repetitive copies of transgene or rDNA. During meiosis (*left*), unpaired DNA can be sensed, resulting in the production of aRNAs. aRNAs are converted to dsRNAs by SAD1, SAD2, and SAD3 in the nuclear periphery. dsRNAs are processed by DCL-1 into sRNAs, which are then loaded onto an SMS-2/QIP-based RISC to accomplish gene silencing. Abbreviations: RNAi, RNA interference; qiRNA, QDE-2-interacting sRNA; aRNA, aberrant RNA; QDE, quelling-defective; dsRNA, double-stranded RNA; SAD-1/2/3, suppressor of ascus dominance 1/2/3; DCL, Dicer-like protein; sRNA, small noncoding RNA; QIP, QDE-2-interacting protein; RISC, RNA-induced silencing complex; SMS, suppressor of meiotic silencing-2.

Dicer, and siRNAs are then loaded onto the RISC. QDE-2 cleaves and removes the passenger strand with the help of QIP to form an active RISC associated with a single-stranded siRNA guide strand that targets complementary mRNA substrates for degradation. Although the mechanism of quelling downstream of dsRNA is well understood, how repetitive DNA sequences trigger the formation of aberrant DNA structures, which are specifically recognized by QDE-3, RPA, and QDE-1, is not known and should be a major focus of future studies.

qiRNA: A Type of sRNA Induced by DNA Damage

Meiotic silencing by unpaired DNA (MSUD): an RNAi-related gene silencing mechanism that occurs in the presence of unpaired homologous DNA sequences during meiosis Quelling triggered by repetitive transgenes occurs in the vegetative *Neurospora* culture under normal growth conditions. Intriguingly, we recently uncovered a class of QDE-2-interacting sRNA, designated qiRNA, that is specifically induced after *Neurospora* is treated with a DNAdamaging agent (61). qiRNAs originate mostly from the rDNA locus, which is the only highly repetitive sequence in the wild-type *Neurospora* genome. These sRNAs are \sim 21–23 nt in length and generally have a 5' uridine. qiRNA biogenesis requires QDE-1, QDE-3, and the Dicers, indicating that qiRNAs are specific sRNA species made by the RNAi machinery upon DNA damage (61). qiRNAs originate from aRNA precursors. In fact, aRNAs ranging in size from 500 bp to 2 kb from the intergenic spacer regions accumulate in the *dcl* double mutant. The production of aRNA is abolished in the *qde-3* mutant, inferring that the RecQ helicase QDE-3 is required for the biogenesis of DNA-damage-induced aRNA. In addition, QDE-1 is essential for aRNA production, but the conventional RNA polymerases (Pol I, Pol II, and Pol III) are not, suggesting that QDE-1 acts as a DdRP to produce the DNA-damage-induced aRNAs.

DNA damage results in cell arrest, decrease of DNA replication, and protein synthesis. In mutants that are defective in qiRNA synthesis, DNA-damage-induced reduction of protein synthesis is blunted (61). QDE-3 plays a role in DNA damage repair (23, 51). Furthermore, *qde-1* and *dcl* mutants also exhibit increased sensitivity to DNA-damaging agents, suggesting that qiRNAs contribute to DNA damage checkpoints by inhibiting protein synthesis (61). In *Arabidopsis*, RNAi components are enriched in the nucleolus, and rDNA-derived siRNAs facilitate heterochromatin formation (83). Furthermore, the *Drosophila* RNAi-deficient mutants have disorganized nucleoli and rDNA loci (82). These studies suggest that rDNA-derived sRNAs have a shared role in eukaryotes to maintain genome integrity.

Both quelling and qiRNA pathways appear to require the same components, suggesting that they are mechanistically similar (60, 61). The repetitive nature of transgenes and the rDNA locus is very likely to be the common trigger for quelling and qiRNA production. The production of qiRNA is induced by DNA damage, implying that replication stress or double-stranded breaks trigger formation of aberrant DNA structures. Although quelling occurs under normal growth conditions, it is likely that the transgene integration loci are fragile DNA sites with elevated levels of replication stress (28, 35). If so, quelling should also be the result of DNA damage at the transgene integration loci trigged by the presence of repetitive DNA sequences.

MEIOTIC SILENCING BY UNPAIRED DNA

Meiotic silencing by unpaired DNA (MSUD) was originally discovered during studies of the *as-cospore maturation 1* gene (*asm-1*) in *Neurospora* (3, 93, 94). Unlike quelling, which takes place in the vegetative stage, MSUD occurs during meiosis. *Neurospora* is a haploid ascomycete and becomes transiently diploid during meiosis following the fusion of two haploid nuclei from opposite mating types. MSUD occurs in prophase I of meiosis when unpaired DNA sequences are present and leads to the silencing of all homologous genes in the diploid ascus cell (94). It has been proposed that *trans*-sensing and meiotic silencing are two related but distinct steps in MSUD (53, 84). The *trans*-sensing step should occur in the nucleus, where the presence of unpaired DNA during meiosis is sensed and likely triggers the production of unpaired DNA-specific aRNA, whereas the meiotic silencing step converts the aRNA to dsRNA and sRNA, resulting in the sRNA-mediated gene silencing. Although, the mechanism of *trans*-sensing is still largely unclear, DNA methylation appears to influence *trans*-sensing without affecting silencing (84). MSUD also operates in the homothallic fungus *Gibberella zeae* (96).

A breakthrough in understanding the mechanism of MSUD was achieved by forward genetic screens for mutants that suppress meiotic silencing (94). The first two genes identified to be required for MSUD are *sad-1* (*suppressor of ascus dominance 1*), which encodes an RdRP and is a paralog of QDE-1 (93, 94), and *sad-2*, which does not contain any conserved domain (5, 95). SAD-1 and SAD-2 colocalize at the perinuclear region and interact physically. A functional *sad-2* gene is required for the proper localization of SAD-1, indicating that SAD-2 might function to recruit SAD-1 to the perinuclear region. The fact that SAD-1 encodes a putative cellular RdRP indicates that MSUD is an RNAi-related phenomenon. In addition, *sms-2* (*suppressor of meiotic silencing 2*), an Argonaute protein homolog in *Neurospora*, is also required for MSUD (59). Although an *sms-2* mutant did not show obvious defects during vegetative growth, the homozygous crosses of *sms-2* mutants are completely barren. In contrast, QDE-2 is not required for MSUD, indicating that there are two parallel RNAi pathways functioning separately in the vegetative and meiotic stages (59).

MSUD was further established as an RNAi-related pathway from studies on the role of *Neurospora* Dicer proteins in meiotic silencing (1). Unlike the single mutants of *sad-1* and *sad-2*, the *dcl-1* single mutants did not function as dominant suppressors of meiotic silencing (1, 94, 95). However, when DCL-1 is inactivated at a later stage of the sexual cycle, meiotic silencing is suppressed, indicating a critical role for DCL-1 in MSUD and therefore a requirement for dsRNA and sRNA in meiotic silencing. In addition, DCL-1 colocalizes with other MSUD components, including SAD-1, SAD-2, and SMS-2, at the nuclear periphery (1, 95). Interestingly, DCL-2, the major Dicer in the quelling pathway, is not required for MSUD (14). QIP, the exonuclease required for RISC activation in the quelling pathway, is also involved in MSUD (42, 60, 109). Like DCL-1, QIP localizes at the perinuclear region with other meiotic silencing components (109).

Recently, a putative RNA/DNA helicase, SAD-3, was demonstrated to be required for meiotic silencing (42). SAD-3 is homologous to *S. pombe* Hrr1, which is a component of the RNA-directed RNA polymerase complex required for the RNAi-mediated heterochromatin formation in fission yeast (75). This link raises the possibility that MSUD and the RNAi-induced heterochromatin assembly may be mechanistically related. SAD-3 is also found in the perinuclear region and associates with SAD-1, SAD-2, and SMS-2 (42). The fact that all the known components involved in MSUD are localized at the perinuclear region and associate with each other suggests that they form a large meiotic silencing complex in this region. It is likely that this complex functions by converting the aRNA produced in the nucleus to dsRNA and then sRNA to trigger gene silencing.

A simple model of the MSUD pathway is shown in **Figure 1**. During meiosis, the pairing of the homologous chromosomes leads to the detection of the unpaired DNA region and production of aRNA transcripts. In the perinuclear region, aRNA transcripts are converted to dsRNA by SAD-1, with the help of SAD-2 and SAD-3. DCL-1 processes dsRNA into sRNAs, which are then loaded onto an SMS-2/QIP-based RISC to execute posttranscriptional silencing of homologous genes. Despite the identification of many components involved in the meiotic silencing step of the MSUD pathway, little is known about the *trans*-sensing mechanism. First, what is the mechanism that scans and senses the unpaired DNA in homologous chromosomes during meiosis? DNA methylation is known to be involved at this step (84), suggesting that chromatin structures are important. Second, what is the DdRP that produces the aRNA and how is the specificity achieved? In quelling, QDE-1 serves as both DdRP and RdRP (60). It will be interesting to test whether QDE-1/SAD-1 has a similar role in MSUD. In addition, some genes appear to be immune to MSUD, such as rDNA clusters and the mating-type genes *mat* A and *mat* a, which have very different DNA sequences (92, 94). Thus, there must be a mechanism that protects certain genes from silencing during meiosis in *Neurospora*. Finally, since the discovery of meiotic silencing in *Neurospora*, unpaired homologous

RNA-induced transcriptional silencing (RITS):

an RNAi-related mechanism that triggers or maintains the formation of heterochromatin, resulting in the silencing of gene expression at the level of transcription DNA-triggered silencing during meiosis in *C. elegans, D. melanogaster*, and mammals has been described (53). Although the silencing mechanisms in *D. melanogaster* and mammals appear to be independent of RNAi, it is still not clear whether MSUD in *Neurospora* is mechanistically similar to the silencing phenomena in animals.

RNAi-MEDIATED HETEROCHROMATIN FORMATION IN SCHIZOSACCHAROMYCES POMBE

Heterochromatin was initially considered to be transcriptionally inert. Heterochromatin in the fission yeast *S. pombe* is distributed in three different loci: telomeres, centromeres, and mating-type loci. The centromeres of fission yeast are similar to those of humans in their structure and epigenetic modifications. The central core region sequence is unique for every centromere, and it is at this region where the kinetochore binds. It is flanked by two types of repeated DNA sequences, the innermost (*imr*) and outermost (*otr*) repeats. The outermost pericentromeric repeats are further composed of the dg and dh sequences; these sequences are coated with histone H3 that is methylated at lysine 9 (H3K9). These specialized chromatin modifications provide docking sites for chromodomain proteins that maintain the transcriptionally silent status of the heterochromatin (29).

A direct link between heterochromatin formation and an RNAi pathway in the fission yeast was first established in 2002, when Volpe et al. (105) showed that RNAi components are required for heterochromatin formation in the centromeric regions. In addition, transcription is detected in the heterochromatin region and the heterochromatic RNAs accumulate in mutants of the RNAi pathway, including *RNA-dependent RNA polymerase (rdp1), dicer (dcr1)*, and *argonaute (ago1)*. The transcripts arising from the heterochromatin regions are mapped to both strands of DNA, indicating a dsRNA precursor. Reduced H3K9 methylation and increased H3K4 methylation are also observed in these mutants. Furthermore, sRNAs derived from the centromeric region are detected, and Rdp1 is enriched at the centromere (105). Together, these results demonstrated the requirement of the *S. pombe* RNAi pathway in heterochromatin formation.

In fission yeast, heterochromatin at the centromeric regions is formed through a self-enforcing loop orchestrated by the RNAi components and chromatin-modifying components (7, 37, 65, 73, 103, 104) (**Figure 2**). This self-enforcing loop is brought about by the transcription of the pericentromeric nascent transcripts by RNA Pol II during the S phase of the cell cycle. Rdp1 converts the nascent transcripts to dsRNA, which Dcr1 recognizes and processes into siRNA. The siRNA is then loaded onto Ago1 to form the RNA-induced transcriptional silencing (RITS) complex (101, 102).

RNA Pol II appears to be the DdRP responsible for the production of centromeric siRNA in *S. pombe*. The role of Pol II in this process was supported by the identification of two point mutations in the RNA Pol II subunits (Rpb2 and Rpb7) (52, 114). These mutations resulted in reduced levels of heterochromatic marks and siRNA production. Genome-wide transcriptional analyses suggest that these defects are not due to an indirect effect caused by changes in global gene expression. Furthermore, neither mutation altered the association of Pol II with the centromeric DNA repeats, suggesting that the defects in siRNA production in these mutants are not due to impaired recruitment of Pol II to centromeric DNA.

The identification of the RITS complex provided a mechanistic link between siRNA and heterochromatin assembly (101). The RITS complex consists of Ago1, Chp1, and Tas3. Chp1 is a chromodomain-containing protein that binds the histone H3 lysine 9 methylation (H3K9me), which recruits the RITS to the pericentromeric region. The active Ago1 bound to a single-stranded



Figure 2

Model for RNAi-mediated heterochromatin formation in *Schizosaccharomyces pombe*. The siRNA-guided RITS complex targets the centromeric nascent transcript synthesized by RNA Pol II during the S phase of the cell cycle. RITS associates with the RDRC complex, where the RdRP activity of the RDRC complex generates more dsRNA. Dcr1 processes the dsRNA into siRNA, feeding into the loop. The CLRC complex associates with the RITS complex, which comes in close proximity to chromatin. Clr4 methylates H3K9 and allows Swi6 (HP1 homolog) to dock at the modified chromatin, thereby forming heterochromatin. Abbreviations: RNAi, RNA interference; siRNA, small interfering RNA; RITS, RNA-induced transcriptional silencing; Pol, polymerase; RDRC, RNA-dependent RNA polymerase complex; dtRP, RNA-dependent RNA polymerase; dsRNA, double-stranded RNA; Dcr1, Dicer 1; CLRC, Clr4-Rik1-Cul4 complex; otr, outer most repeat; imr, inner most repeat; Ago1, Argonaute 1; Tas3, tyrosine auxotrophy suppressor 3; Clr4, calcitonin-like receptor 4; Chp1, chromodomain-containing protein 1; Swi6, SWItch/sucrose nonfermentable 6; Rik1, RS2-interacting KH domain protein 1; Dos1/2, delocalization of Swi6 1/2; Stc1, siRNA to chromatin; Rdp1, RNA-dependent RNA polymerase 1; Hrr1, helicase required for RNAi-mediated heterochromatin assembly; Cid12, caffeine-induced death resistant 12.

guide siRNA also reinforces this interaction through homology-dependent association with the centromeric nascent RNA transcripts.

The production of the nascent transcripts provides a template for Rdp1 (75). Rdp1 is part of the RNA-dependent RNA polymerase complex (RDRC) that also contains Hrr1 and Cid12 (75). Both Rdp1 and Hrr1 associate with heterochromatin and centromeric (*cen*) RNA transcripts (*cen* RNA). Rdp1 generates dsRNA in a primer-independent manner from single-stranded *cen* RNA, and it has been proposed that the RITS complex acts as a priming complex for RDRC to localize at the centromeric DNA. Dicer cleaves the dsRNA into siRNA and greatly enhances the activity of RDRC in siRNA production in vitro (24). Although Dcr1 is present mostly in the cytoplasm (43), a study showed that Dicer is also present in the nucleus, suggesting that Dcr1-mediated siRNA production occurs in *cis* with transcription (30). Dcr1 may shuttle between the nucleus and cytoplasm, and the retention of Dcr1 in the nucleus is essential for the production of centromeric siRNAs.

The recruitment of RDRC to the centromeric region is dependent on Clr4 (75). Clr4, a histone methyltransferase, is recruited to the pericentromeric region, where it methylates H3K9. The methylated chromatin further recruits Clr4 and the chromodomain proteins Swi6, Chp1, and Chp2 to the region (86, 111). siRNA production is greatly reduced in the *clr4* mutants. Clr4

associates with Rik1, Dos1, Dos2, and Cul4 in a complex called Clr4-Rik1-Cul4 complex (CLRC); all the components in this complex are required for heterochromatin assembly (44, 45, 48, 66). The association of Swi6 to the heterochromatin allows the recruitment of cohesion, whereas the binding of Clr4 and Chp1 stabilizes CLRC and RITS at the heterochromatin. Together, this allows the reinforcement of silencing signals and spreading of heterochromatin (**Figure 2**).

Two models have been proposed to explain how RITS is recruited to specific chromosome regions. In the first model, RITS is guided by siRNA to the centromeric DNA through direct base-pairing, which would require unwinding of the DNA double helix. In the second model, the siRNA is targeted to the nascent transcript produced from the pericentromeric region. RNA immunoprecipitation demonstrated that RITS and RDRC are associated with the noncoding *cen* RNA and that this association requires Clr4 methyltransferase and Dcr1 (75). These results suggest that recruitment of RITS to the centromeric regions is not direct but rather is mediated through the siRNA-targeted binding of AGO1 to the nascent transcripts (7, 8, 107). This RNA platform model was tested in vivo by tethering Tas3 (a component of the RITS complex) to the RNA led to transcriptional silencing of *ura4*⁺ (4, 7). This study demonstrates that the association of RITS with the nascent transcript is sufficient to trigger H3K9me and transcriptional gene silencing.

The formation of heterochromatin in the centrometric regions is cell cycle dependent (16, 56). The DNA replication of the heterochromatin occurs in early S phase. This is also precisely the time during the cell cycle when pericentromeric dg/db transcripts are detected. In addition, Pol II localization to the centromeric regions is detected in early S phase when the level of H3K9 methylation is at its lowest level. In S phase, RNAi components accumulate at the pericentromeric region and H3K9 methylation marks are detected. By the end of replication, the epigenetic marks of heterochromatin are faithfully passed on to the sister chromatids. The H3K9me levels further increase as the cell cycle progresses and peak during the G2 phase of the cell cycle. Thus, the RNAi components in fission yeast aid in the formation of heterochromatin and assist the cell in completing DNA synthesis in heterochromatic regions (110). Cotranscriptional RNAi releases the RNA Pol II from the centromeric regions, which allows the completion of DNA replication and subsequent spreading of histone modifications.

Recent advances in our understanding of the RNAi-mediated heterochromatin formation have demonstrated how RNAi and chromatin-modifying components are recruited specifically to the centromeric loci, and have revealed the time frame in which these processes occur. However, several recent studies have raised questions regarding the initial trigger that results in heterochromatin formation at the centromeric regions. Halic & Moazed (39) proposed that the production of primal RNAs (priRNAs) are initiators of this process. These priRNAs are thought to be the products of degraded transcripts, are generated independently of Dcr1 and RDRC, and are loaded onto Ago1. The primed Ago1-priRNAs complexes target the centromeric loci and further recruit Clr4, which methylates H3K9 and initiates the formation of RNAi-mediated heterochromatin. Results of another study led to the alternative hypothesis that the CLRC complex is recruited to the centromeric regions independently of RNAi components (91). Consistent with this hypothesis, overexpression of Clr4 in a clr4 ago1 double mutant induced de novo H3K9me2 at the centromeric regions, suggesting RNAi may be dispensable for initiating heterochromatin formation but critical in maintaining the heterochromatic status. Forcibly tethering the Clr4 methyltransferase to a euchromatic region is also sufficient to establish heterochromatin formation in the absence of RNAi components (50). Although a tremendous amount of progress has been made, further studies are needed to fully understand the mechanism of RNAi-mediated heterochromatin formation in the fission yeast.

RNAi IN BUDDING YEASTS

Although the RNA-silencing pathway is highly conserved throughout the fungal kingdom, some budding yeasts such as *Saccharomyces cerevisiae* lack homologs of Argonaute, Dicer, and RdRP. However, sequence homologs of Argonaute but not other RNAi components are present in some of the budding yeasts, such as *Saccharomyces castellii*, *Kluyveromyces polysporus*, and *Candida albicans* (27). These yeasts use noncanonical Dicer proteins to produce siRNAs, which map mostly to transposable elements and subtelomeric repeats. Remarkably, the expression of the *S. castellii argonaute* and *dicer* genes in *S. cerevisiae* activates an RNAi pathway that inhibits transposon transposition. These results suggest that the RNAi pathway is an ancient host defense mechanism that has been lost in some fungal species.

miRNA-LIKE sRNA IN NEUROSPORA

miRNAs are endogenous ~21-nt RNA duplexes processed by Dicers from ssRNA precursors with a hairpin structure. miRNAs in animals, plants, and algae have been discovered and regulate physiological and developmental processes by targeting mRNA for cleavage or translational repression (2, 6, 38, 57, 63, 64, 69, 74). miRNA was thought to be absent in fungi until our recent identification of miRNA-like sRNAs (milRNAs) in *Neurospora* (62). At least 25 *Neurospora* milRNA-producing loci have been identified. Several lines of evidence suggest that milRNAs in *Neurospora* regulate gene expression similar to animal and plant miRNAs. First, the mRNA of predicated target genes of *milR-1* are upregulated in the *dcl* and *qde-2* mutants. Second, the expression of a reporter gene containing an *milR-1* target site is increased in the *qde-2* strain. Third, QDE-2 specifically associates with the predicted *milR-1* target mRNAs. Interestingly, milRNAs do not appear to play critical roles in *Neurospora* growth or developmental processes because *qde-2* or milRNA knock-out strains grow normally. Future studies are needed to establish the physiological importance of milRNAs in *Neurospora*. Although currently no miRNA-like sRNA has been experimentally confirmed in other fungal species, a study of the wheat pathogen *Mycosphaerella graminicola* found 65 potential milRNA-producing loci that produce sRNAs (36).

Unlike the animal and plant miRNAs, the *Neurospora* milRNAs are produced by at least four different biogenesis pathways that utilize different combinations of components, including Dicers, QDE-2, QIP, and an RNase III domain-containing protein, MRPL3 (62). Of the four miRNAs that were investigated in detail, only the biogenesis of *milR-3* is mechanistically similar to that of miRNAs in plants; it requires only Dicer for the generation of pre-milRNA and milRNA (49). The production of *milR-4* is only partially dependent on Dicers, indicating the involvement of other nucleases in milRNA biogenesis. *milR-1* is the most abundant sRNA-producing locus in the *Neurospora* genome, and production of milR-1 requires Dicers, QDE-2, and catalytically active QIP. The primary (pri)-*milR-1* is first processed by Dicer to produce pre-*milR-1*. Afterward, QDE-2 binds to pre-*milR-1* and recruits the exonuclease QIP to process the pre-*milR-1* into mature *milR-1*. In this pathway, the Argonaute QDE-2 functions as a scaffold for *milR-1* maturation.

Although biogenesis of *milR-1*, *milR-3*, and *milR-4* are completely or partially dependent on Dicers, the biogenesis of *milR-2* is completely independent of Dicer. Instead, the production of *milR-2* requires catalytically active QDE-2. Our results suggest that QDE-2 binds to the premil-RNA and cleaves the passenger strand using its slicer activity. The *milR-2* biogenesis pathway is the first known example of a Dicer-independent—but Argonaute-dependent—mechanism for sRNA production. Soon after our study of *milR-2* biogenesis, *miR-451* in both mouse and zebrafish was shown to be produced by a similar mechanism (15, 19), suggesting that this is a conserved sRNA biogenesis pathway. The surprisingly diverse milRNA pathways in *Neurospora* shed important light on the biogenesis and evolution of eukaryotic miRNAs.

milRNA: microRNA-like small RNA

RNAI AS A HOST DEFENSE MECHANISM IN FUNGI

Transposon Control in Neurospora

Quelling results in the silencing of repetitive transgenes in *Neurospora*, suggesting that quelling is a mechanism that silences transposons. In most *Neurospora* strains, there are no functional transposons because of repeat-induced point mutation, a process that can mutate repetitive sequences during meiosis (33). However, a functional long interspersed element (LINE)-like transposon was previously identified in an African strain (55), and Nolan et al. (77) introduced the Tad transposon into laboratory *Neurospora* strains and showed that QDE-2 and Dicer, but not QDE-1 and QDE-3, are required for the suppression of transposon replication. These results suggest that transposition may generate inverted repeats that lead to the production of dsRNA, resulting in the silencing of transposon.

An Antiviral Mechanism in Cryphonectria parasitica and Aspergillus nidulans

Virus infection often leads to the production of siRNAs derived from viral dsRNA, suggesting a role for RNAi as part of the innate immunity mechanism against viral replication (26). Plants, arthropods, nematodes, and some other animals rely on the RNAi pathway to protect them from viral infection. To counter the effects of RNAi, viruses often express viral suppressors of RNA silencing that can inhibit RNAi. In fungi, an RNAi-based antiviral mechanism was first observed in the ascomycete filamentous fungus *Cryphonectria parasitica* (90, 97, 112, 113). *C. parasitica* is a rich resource for the study of basic aspects of virus-fungus interactions because it can support the replication of five virus families and is a well-established experimental system for the study of mycoviruses.

Like *Neurospora*, *C. parasitica* has two *dicer*-like genes, *dcl1* and *dcl2*. Hypovirus or mycoreovirus infection of the *dcl2* mutant results in a severe growth phenotype and elevated viral RNA levels (90), indicating that DCL2 is the major Dicer in *C. parasitica*. Consistently, 21- to 22-nt virus-derived sRNAs were abolished in the *dcl2* mutant (113). Although there are four Argonaute-like genes in *C. parasitica*, only *agl2* is required for the antiviral defense response (97). Together, these observations demonstrate that RNAi acts as a defense mechanism against viruses.

The mycovirus *Cryphonectria* hypovirus 1 (CHV1) expresses p29, a papain-like protease similar to the plant potyvirus-encoded suppressor of RNA silencing HC-Pro (17, 89, 98, 99). Deletion of p29 results in the reduction of viral RNA levels. p29 suppresses hairpin RNA-induced, virus-induced, and agroinfiltration-induced RNA silencing, indicating that it is a suppressor of RNA silencing. When *C. parasitica* is infected by a mutant hypovirus without p29, the mRNAs of *dcl*2 and *agl2* accumulate to high levels, suggesting that p29 represses RNAi by inhibiting the expression of RNA igenes.

In addition to their roles in silencing viral replication by degrading viral RNAs, RNAi components also promote viral RNA recombination in *C. parasitica*, which leads to the production of hypovirus defective-interfering (DI) RNAs (97, 112). DI RNAs are a result of deletion events during recombination; DI RNAs suppress parental RNA accumulation, causing symptoms to attenuate. Importantly, DI RNAs do not accumulate in *agl1*, *agl2*, or *dcl2* mutants, indicating that these RNAi components are also involved in hypoviral RNA recombination and DI RNA production. The RNAi components may contribute to viral recombination by providing 5' and 3' fragments of viral RNA to the viral RdRP, but studies are needed to validate such a mechanism.

Studies of the RNAi pathway in *Aspergillus nidulans* also provided insights into the fungusvirus interaction. *A. nidulans* has one Argonaute gene and one Dicer gene; both are required for dsRNA-triggered gene silencing in the organism (41). Although there are two RdRP genes in the genome, neither is required for dsRNA-induced gene silencing, suggesting the absence of an sRNA amplification step by RdRPs (41). Infection with the *Aspergillus* virus 1816 suppresses dsRNA-induced RNA silencing (40), indicating the existence of an RNA-silencing suppressor encoded by this virus. In addition, the virus 341–derived siRNA was detected at a high level in the Argonaute mutant, indicating that this virus is targeted by the RNAi machinery (40).

dsRNA/Viral Infection–Induced Transcriptional Response in *Neurospora* and *C. parasitica*

Innate immunity, triggered after an organism detects dsRNA or foreign particles, is an important part of the eukaryotic host defense mechanism. The mechanism activates a signaling cascade, which in turn activates the transcription of effector genes that defend against the foreign invader. In *Neurospora*, a similar transcriptional response is observed upon the induction of dsRNA expression, which induces the expression of key RNAi components, QDE-2 and DCL2, and other putative antiviral genes are upregulated (18). The transcriptional response is retained in the *dcl1 dcl2* double mutant, indicating that the response is induced by dsRNA, not siRNA. Furthermore, the known RNAi components in *Neurospora* are not required for this response. Interestingly, dsRNA significantly induces the expression of QDE2 both transcriptionally and posttranscriptionally. In the *dcl* double mutant, despite the induction of *qde-2* transcription, the QDE-2 protein levels stay constant after dsRNA expression. This suggests that siRNAs are important for stabilizing the QDE2 protein. The induction of QDE-2 by dsRNA is required for efficient RNAi because mutants that lack this response have lower RNAi efficiency.

Genome-wide analyses identified approximately 60 genes that are activated by dsRNA. In addition to RNAi components, many of these genes are homologs of antiviral and interferon-stimulated genes, suggesting that the dsRNA-induced transcriptional response is part of an ancient host defense response. Because *Neurospora* does not encode homologs of the known mammalian dsRNA sensors, the transcriptional program triggered by dsRNA should represent a novel signaling cascade.

Interestingly, in *C. parasitica*, the expression of *dcl2* and *agl2* transcripts is strongly induced upon viral infection and expression of hairpin RNA (112, 113), indicating that this antiviral transcriptional activation response is conserved in filamentous fungi. In addition, the deletion of p29, also results in upregulation of both *dcl2* and *agl2*. Moreover, by using a GFP reporter fused with the promoter of *agl2*, Sun et al. (97) demonstrated that the induction of *agl2* by viral infection is regulated at the transcriptional level. Interestingly, the viral infection–induced *dcl-2* expression is blocked in the *agl2* mutant, suggesting that AGL2 plays a regulatory role in this gene activation pathway. These results suggest that the transcriptional regulation of RNAi components is important for the host antiviral response.

RNA SILENCING AND SMALL RNA STUDIES ON OTHER FILAMENTOUS FUNGI

Gene Silencing and Small RNAs in Mucor circinelloides

Mucor circinelloides, which causes invasive maxillofacial zygomycosis, is an important model system for studying RNAi in the zygomycete clade. Transformation with hairpin RNA-producing constructs and self-replicative plasmids shows that this organism can silence gene expression. Silencing results in the accumulation of two size classes of siRNAs (21 and 25 nt). *M. circinelloides* has two redundant *dicer (dcl1* and *dcl2)* genes and two *rdrp* genes (*rdrp1* and *rdrp2*); DCL-2 plays

the major role in the RNA-silencing pathway and is induced by both sense and hairpin transgenes. Secondary sRNA were detected, suggesting an sRNA amplification step in the RNA-silencing pathway.

Detailed examination of the sRNA profile in this fungus revealed four classes of endogenous small RNA (esRNA). Interestingly, the esRNAs detected in *Mucor* are derived from exons and can regulate mRNA accumulation. The largest class of these exon-derived siRNAs (ex-siRNA) is dependent on DCL-2 and RDRP1, suggesting that mRNAs from these loci are converted to dsRNA by RdRP1 and processed by DCL-2. The second class of ex-siRNAs requires DCL-2 and RdRP2 for biogenesis. The third class requires both RdRP1 and RdRP2, and the two Dicers have redundant roles in their biogenesis. The fourth class requires DCL1, but not DCL2, and the two RdRPs are required. These results indicate that different RNAi components are used to produce different ex-siRNAs.

RNAi Studies on the Human Fungal Pathogen Cryptococcus neoformans

Cryptococcus neoformans, which causes fatal meningoencephalitis, has Argonaute, Dicer, and RdRP genes (47, 67). The RNAi pathway in *C. neoformans* acts in a process called sex-induced silencing (SIS) (106). SIS, which is similar to quelling, requires multiple copies of transgenes. In addition, the efficiency of SIS increases dramatically during the sexual cycle, which is correlated with increased expression of known RNAi components. SIS and SIS-associated sRNA production is abolished in RNAi mutants, indicating the role of RNAi in this silencing process during the sexual cycle. Examination of the sRNA profile revealed that some of the sRNAs, which were not derived from the transgenes, mapped to repetitive transposable elements and presumptive centromeric regions, suggesting that the RNAi pathway is a genome defense mechanism. Supporting this notion, in the *rdp1* mutant, retrotransposons are highly expressed with higher transposition rates.

Small RNAs from Magnaporthe oryzae

Magnaporthe oryzae is a model organism for the study of pathogen-host interactions because it is a primary plant pathogen of rice and wheat. A recent sRNA analysis revealed that a large variety of sRNAs map to loci, including tRNA loci, rRNA loci, coding regions, and intergenic loci (79). In the mycelia, sRNAs mostly come from intergenic regions and repetitive elements, such as long terminal repeat retrotransposons. However, sRNAs in the appressoria are enriched for tRNAderived RNA fragments. The differential distribution of sRNAs in different tissues suggests these sRNAs have roles in the regulation of growth and development of this pathogen.

SUMMARY POINTS

- 1. *Neurospora crassa* has multiple RNAi-related silencing pathways, including quelling, qiRNA, and MSUD, which are triggered by multiple copies of transgenes, DNA damage, and unpaired DNA, respectively.
- 2. Schizosaccharomyces pombe requires an RNAi pathway for heterochromatin formation.
- 3. There are multiple sRNA pathways in fungi. milRNAs with distinct biogenesis pathways that require combinations of different components have been found in *Neurospora*.
- 4. RNAi is an important fungal host defense mechanism against transposon and viral invasion.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

We thank the members of the Liu laboratory for discussion and comments. This work was supported by grants from the National Institutes of Health and the Welch Foundation to Yi Liu (I-1560).

LITERATURE CITED

- 1. Alexander WG, Raju NB, Xiao H, Hammond TM, Perdue TD, et al. 2008. DCL-1 colocalizes with other components of the MSUD machinery and is required for silencing. *Fungal Genet. Biol.* 45:719–27
- Ambros V, Bartel B, Bartel DP, Burge CB, Carrington JC, et al. 2003. A uniform system for microRNA annotation. RNA 9:277–79
- 3. Aramayo R, Metzenberg RL. 1996. Meiotic transvection in fungi. Cell 86:103-13
- Bühler M, Verdel A, Moazed D. 2006. Tethering RITS to a nascent transcript initiates RNAiand heterochromatin-dependent gene silencing. *Cell* 125:873–86
- Bardiya N, Alexander WG, Perdue TD, Barry EG, Metzenberg RL, et al. 2008. Characterization of interactions between and among components of the meiotic silencing by unpaired DNA machinery in *Neurospora crassa* using bimolecular fluorescence complementation. *Genetics* 178:593–96
- 6. Bartel DP. 2004. MicroRNAs: genomics, biogenesis, mechanism, and function. Cell 116:281-97
- Buhler M, Moazed D. 2007. Transcription and RNAi in heterochromatic gene silencing. Nat. Struct. Mol. Biol. 14:1041–48
- Cam HP, Chen ES, Grewal SIS. 2009. Transcriptional scaffolds for heterochromatin assembly. *Cell* 136:610–14
- Carattoli A, Cogoni C, Morelli G, Macino G. 1994. Molecular characterization of upstream regulatory sequences controlling the photoinduced expression of the *albino-3* gene of *Neurospora crassa*. *Mol. Microbiol.* 13:787–95
- Carthew RW, Sontheimer EJ. 2009. Origins and mechanisms of miRNAs and siRNAs. Cell 136:642– 55
- 11. Catalanotto C, Azzalin G, Macino G, Cogoni C. 2000. Gene silencing in worms and fungi. *Nature* 404:245
- 12. Catalanotto C, Azzalin G, Macino G, Cogoni C. 2002. Involvement of small RNAs and role of the *qde* genes in the gene silencing pathway in *Neurospora*. *Genes Dev*. 16:790–95
- Catalanotto C, Nolan T, Cogoni C. 2006. Homology effects in Neurospora crassa. FEMS Microbiol. Lett. 254:182–89
- 14. Catalanotto C, Pallotta M, ReFalo P, Sachs MS, Vayssie L, et al. 2004. Redundancy of the two Dicer genes in transgene-induced posttranscriptional gene silencing in *Neurospora crassa*. *Mol. Cell. Biol.* 24:2536–45
- Cheloufi S, Dos Santos CO, Chong MM, Hannon GJ. 2010. A Dicer-independent miRNA biogenesis pathway that requires Ago catalysis. *Nature* 465:584–89
- Chen ES, Zhang K, Nicolas E, Cam HP, Zofall M, Grewal SIS. 2008. Cell cycle control of centromeric repeat transcription and heterochromatin assembly. *Nature* 451:734–37
- Choi GH, Pawlyk DM, Nuss DL. 1991. The autocatalytic protease p29 encoded by a hypovirulenceassociated virus of the chestnut blight fungus resembles the potyvirus-encoded protease HC-Pro. *Virology* 183:747–52
- Choudhary S, Lee H-C, Maiti M, He Q, Cheng P, et al. 2007. A double-stranded-RNA response program important for RNA interference efficiency. *Mol. Cell. Biol.* 27:3995–4005
- Cifuentes D, Xue H, Taylor DW, Patnode H, Mishima Y, et al. 2010. A novel miRNA processing pathway independent of Dicer requires Argonaute2 catalytic activity. *Science* 328:1694–98

4. Demonstrates that the targeting of RITS complex via RNA to a chromosomal locus is sufficient to induce transcriptional gene silencing.

11. Identifies the Argonaute protein QDE-2 as a factor required for quelling, indicating that dsRNA-induced gene silencing in animals and quelling in *Neurospora* share a common mechanism.

16. Discovered that siRNA production and heterochromatin formation are regulated by the cell cycle and suggested a role for RNAi in the cell cycle.

Annu. Rev. Microbiol. 2012.66:305-323. Downloaded from www.annualreviews.org by Universidade Federal do Parana on 03/05/13. For personal use only.

- 22. Identified the first eukaryotic RNAi component and suggested that dsRNA is an endogenous trigger for posttranscriptional gene silencing.
- Cogoni C, Irelan JT, Schumacher M, Schmidhauser TJ, Selker EU, Macino G. 1996. Transgene silencing of the *al-1* gene in vegetative cells of *Neurospora* is mediated by a cytoplasmic effector and does not depend on DNA-DNA interactions or DNA methylation. *EMBO 7.* 15:3153–63
- Cogoni C, Macino G. 1997. Isolation of quelling-defective (qde) mutants impaired in posttranscriptional transgene-induced gene silencing in Neurospora crassa. Proc. Natl. Acad. Sci. USA 94:10233–38
- 22. Cogoni C, Macino G. 1999. Gene silencing in *Neurospora crassa* requires a protein homologous to RNA-dependent RNA polymerase. *Nature* 399:166–69
- Cogoni C, Macino G. 1999. Posttranscriptional gene silencing in *Neurospora* by a RecQ DNA helicase. Science 286:2342–44
- Colmenares SU, Buker SM, Buhler M, Dlakić M, Moazed D. 2007. Coupling of double-stranded RNA synthesis and siRNA generation in fission yeast RNAi. *Mol. Cell* 27:449–61
- Dang Y, Yang Q, Xue Z, Liu Y. 2011. RNA interference in fungi: pathways, functions and applications. *Eukaryot. Cell* 10:1148–55
- Ding SW, Lu R. 2011. Virus-derived siRNAs and piRNAs in immunity and pathogenesis. Curr. Opin. Virol. 1:533–44
- Drinnenberg IA, Weinberg DE, Xie KT, Mower JP, Wolfe KH, et al. 2009. RNAi in budding yeast. Science 326:544–50
- 28. Durkin SG, Glover TW. 2007. Chromosome fragile sites. Annu. Rev. Genet. 41:169-92
- 29. Ebert A, Lein S, Schotta G, Reuter G. 2006. Histone modification and the control of heterochromatic gene silencing in *Drosophila. Chromosome Res.* 14:377–92
- Emmerth S, Schober H, Gaidatzis D, Roloff T, Jacobeit K, Bühler M. 2010. Nuclear retention of fission yeast Dicer is a prerequisite for RNAi-mediated heterochromatin assembly. *Dev. Cell* 18:102–13
- Fire A, Xu S, Montgomery MK, Kostas SA, Driver SE, Mello CC. 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* 391:806–11
- Fulci V, Macino G. 2007. Quelling: post-transcriptional gene silencing guided by small RNAs in Neurospora crassa. Curr. Opin. Microbiol. 10:199–203
- 33. Galagan JE, Selker EU. 2004. RIP: the evolutionary cost of genome defense. Trends Genet. 20:417-23
- Ghildiyal M, Xu J, Seitz H, Weng Z, Zamore PD. 2010. Sorting of *Drosophila* small silencing RNAs partitions microRNA* strands into the RNA interference pathway. *RNA* 16:43–56
- 35. Glover TW. 2006. Common fragile sites. Cancer Lett. 232:4-12
- 36. Goodwin SB, Ben M'Barek S, Dhillon B, Wittenberg AHJ, Crane CF, et al. 2011. Finished genome of the fungal wheat pathogen *Mycosphaerella graminicola* reveals dispensome structure, chromosome plasticity, and stealth pathogenesis. *PLoS Genet.* 7:e1002070
- Grewal SI. 2010. RNAi-dependent formation of heterochromatin and its diverse functions. Curr. Opin. Genet. Dev. 20:134–41
- Grimson A, Srivastava M, Fahey B, Woodcroft BJ, Chiang HR, et al. 2008. Early origins and evolution of microRNAs and Piwi-interacting RNAs in animals. *Nature* 455:1193–97
- Halic M, Moazed D. 2010. Dicer-independent primal RNAs trigger RNAi and heterochromatin formation. *Cell* 140:504–16
- Hammond TM, Andrewski MD, Roossinck MJ, Keller NP. 2008. Aspergillus mycoviruses are targets and suppressors of RNA silencing. Eukaryot. Cell 7:350–57
- Hammond TM, Keller NP. 2005. RNA silencing in *Aspergillus nidulans* is independent of RNAdependent RNA polymerases. *Genetics* 169:607–17
- 42. Hammond TM, Xiao H, Boone EC, Perdue TD, Pukkila PJ, Shiu PK. 2011. SAD-3, a putative helicase required for meiotic silencing by unpaired DNA, interacts with other components of the silencing machinery. *Genes Genomes Genet*. 1:369–76
- Hayashi A, Da-Qiao D, Tsutsumi C, Chikashige Y, Masuda H, et al. 2009. Localization of gene products using a chromosomally tagged GFP-fusion library in the fission yeast *Schizosaccharomyces pombe*. *Genes Cells* 14:217–25
- 44. Hong E-JE, Villén J, Moazed D. 2005. A cullin E3 ubiquitin ligase complex associates with Rik1 and the Clr4 histone H3-K9 methyltransferase and is required for RNAi-mediated heterochromatin formation. *RNA Biol.* 2:106–11

- 45. Horn PJ, Bastie J-N, Peterson CL. 2005. A Rik1-associated, cullin-dependent E3 ubiquitin ligase is essential for heterochromatin formation. *Genes Dev.* 19:1705–14
- 46. Hsieh J, Fire A. 2000. Recognition and silencing of repeated DNA. Annu. Rev. Genet. 34:187-204
- Idnurm A, Bahn Y-S, Nielsen K, Lin X, Fraser JA, Heitman J. 2005. Deciphering the model pathogenic fungus Cryptococcus neoformans. Nat. Rev. Micro 3:753–64
- Jia S, Noma K-I, Grewal SIS. 2004. RNAi-independent heterochromatin nucleation by the stressactivated ATF/CREB family proteins. *Science* 304:1971–76
- Jones-Rhoades MW, Bartel DP, Bartel B. 2006. MicroRNAs and their regulatory roles in plants. Annu. Rev. Plant Biol. 57:19–53
- Kagansky A, Folco HD, Almeida R, Pidoux AL, Boukaba A, et al. 2009. Synthetic heterochromatin bypasses RNAi and centromeric repeats to establish functional centromeres. *Science* 324:1716–19
- Kato A, Akamatsu Y, Sakuraba Y, Inoue H. 2004. The *Neurospora crassa mus-19* gene is identical to the *qde-3* gene, which encodes a RecQ homologue and is involved in recombination repair and postreplication repair. *Curr. Genet.* 45:37–44
- 52. Kato H, Goto DB, Martienssen RA, Urano T, Furukawa K, Murakami Y. 2005. RNA polymerase II is required for RNAi-dependent heterochromatin assembly. *Science* 309:467–69
- 53. Kelly WG, Aramayo R. 2007. Meiotic silencing and the epigenetics of sex. Chromosome Res. 15:633-51
- Kim VN, Han J, Siomi MC. 2009. Biogenesis of small RNAs in animals. Nat. Rev. Mol. Cell Biol. 10:126– 39
- Kinsey JA. 1990. Tad, a LINE-like transposable element of *Neurospora*, can transpose between nuclei in heterokaryons. *Genetics* 126:317–23
- Kloc A, Zaratiegui M, Nora E, Martienssen R. 2008. RNA interference guides histone modification during the S phase of chromosomal replication. *Curr. Biol.* 18:490–95
- Lagos-Quintana M, Rauhut R, Lendeckel W, Tuschl T. 2001. Identification of novel genes coding for small expressed RNAs. *Science* 294:853–58
- 58. Lau NC, Seto AG, Kim J, Kuramochi-Miyagawa S, Nakano T, et al. 2006. Characterization of the piRNA complex from rat testes. *Science* 313:363–67
- 59. Lee DW, Pratt RJ, McLaughlin M, Aramayo R. 2003. An Argonaute-like protein is required for meiotic silencing. *Genetics* 164:821–28
- 60. Lee HC, Aalto AP, Yang Q, Chang SS, Huang G, et al. 2010. The DNA/RNA-dependent RNA polymerase QDE-1 generates aberrant RNA and dsRNA for RNAi in a process requiring replication protein A and a DNA helicase. *PLoS Biol.* 8:e1000496
- 61. Lee HC, Chang SS, Choudhary S, Aalto AP, Maiti M, et al. 2009. qiRNA is a new type of small interfering RNA induced by DNA damage. *Nature* 459:274–77
- 62. Lee HC, Li L, Gu W, Xue Z, Crosthwaite SK, et al. 2010. Diverse pathways generate microRNAlike RNAs and Dicer-independent small interfering RNAs in fungi. *Mol. Cell* 38:803–14
- 63. Lee RC, Ambros V. 2001. An extensive class of small RNAs in Caenorhabditis elegans. Science 294:862-64
- Lee RC, Feinbaum RL, Ambros V. 1993. The C. elegans heterochronic gene lin-4 encodes small RNAs with antisense complementarity to lin-14. Cell 75:843–54
- Lejeune E, Bayne EH, Allshire RC. 2010. On the connection between RNAi and heterochromatin at centromeres. Cold Spring Harb. Symp. Quant. Biol. 75:275–83
- 66. Li F, Goto DB, Zaratiegui M, Tang X, Martienssen R, Cande WZ. 2005. Two novel proteins, Dos1 and Dos2, interact with Rik1 to regulate heterochromatic RNA interference and histone modification. *Curr. Biol.* 15:1448–57
- Lin X, Heitman J. 2006. The biology of the Cryptococcus neoformans species complex. Annu. Rev. Microbiol. 60:69–105
- Liu Y, Ye X, Jiang F, Liang C, Chen D, et al. 2009. C3PO, an endoribonuclease that promotes RNAi by facilitating RISC activation. *Science* 325:750–53
- Llave C, Kasschau KD, Rector MA, Carrington JC. 2002. Endogenous and silencing-associated small RNAs in plants. *Plant Cell* 14:1605–19
- Maiti M, Lee HC, Liu Y. 2007. QIP, a putative exonuclease, interacts with the *Neurospora* Argonaute protein and facilitates conversion of duplex siRNA into single strands. *Genes Dev.* 21:590–600

61. Suggests that DNA damage is an important trigger for sRNA production and proposes that the RNA-dependent RNA polymerase also functions to produce aberrant RNAs.

62. Discovered the miRNA-like RNAs in *Neurospora* and uncovered surprisingly diverse biogenesis mechanisms for RNA production.

85. Describe the discovery of quelling, one of the first known RNAi-related phenomena, in *Neurospora*.

75. Discovered the

siRNA production process and

heterochromatin

assembly machinery.

critical link between the

94. Demonstrates that meiotic silencing in *Neurospora* is triggered by unpaired DNA and is an RNAi-related phenomenon.

- Matzke M, Kanno T, Daxinger L, Huettel B, Matzke AJ. 2009. RNA-mediated chromatin-based silencing in plants. Curr. Opin. Cell Biol. 21:367–76
- 72. Mello CC, Conte D. 2004. Revealing the world of RNA interference. Nature 431:338-42
- 73. Moazed D. 2009. Small RNAs in transcriptional gene silencing and genome defence. Nature 457:413-20
- Molnar A, Schwach F, Studholme DJ, Thuenemann EC, Baulcombe DC. 2007. miRNAs control gene expression in the single-cell alga *Chlamydomonas reinbardtii*. *Nature* 447:1126–29
- Motamedi MR, Verdel A, Colmenares SU, Gerber SA, Gygi SP, Moazed D. 2004. Two RNAi complexes, RITS and RDRC, physically interact and localize to noncoding centromeric RNAs. *Cell* 119:789–802
- Napoli C, Lemieux C, Jorgensen R. 1990. Introduction of a chimeric chalcone synthase gene into petunia results in reversible co-suppression of homologous genes in *trans. Plant Cell* 2:279–85
- 77. Nolan T, Braccini L, Azzalin G, De Toni A, Macino G, Cogoni C. 2005. The post-transcriptional gene silencing machinery functions independently of DNA methylation to repress a LINE1-like retrotransposon in *Neurospora crassa*. *Nucleic Acids Res.* 33:1564–73
- Nolan T, Cecere G, Mancone C, Alonzi T, Tripodi M, et al. 2008. The RNA-dependent RNA polymerase essential for post-transcriptional gene silencing in *Neurospora crassa* interacts with replication protein A. *Nucleic Acids Res.* 36:532–38
- Nunes C, Gowda M, Sailsbery J, Xue M, Chen F, et al. 2011. Diverse and tissue-enriched small RNAs in the plant pathogenic fungus, *Magnaporthe oryzae. BMC Genomics* 12:288
- Obbard DJ, Gordon KHJ, Buck AH, Jiggins FM. 2009. The evolution of RNAi as a defence against viruses and transposable elements. *Philos. Trans. R. Soc. B* 364:99–115
- O'Brien HE, Parrent JL, Jackson JA, Moncalvo JM, Vilgalys R. 2005. Fungal community analysis by large-scale sequencing of environmental samples. *Appl. Environ. Microbiol.* 71:5544–50
- Peng JC, Karpen GH. 2007. H3K9 methylation and RNA interference regulate nucleolar organization and repeated DNA stability. *Nat. Cell Biol.* 9:25–35
- Pontes O, Li CF, Costa Nunes P, Haag J, Ream T, et al. 2006. The Arabidopsis chromatin-modifying nuclear siRNA pathway involves a nucleolar RNA processing center. Cell 126:79–92
- Pratt RJ, Lee DW, Aramayo R. 2004. DNA methylation affects meiotic trans-sensing, not meiotic silencing, in *Neurospora. Genetics* 168:1925–35
- 85. Romano N, Macino G. 1992. Quelling: transient inactivation of gene expression in *Neurospora* crassa by transformation with homologous sequences. *Mol. Microbiol.* 6:3343–53
- Sadaie M, Iida T, Urano T, Nakayama J-I. 2004. A chromodomain protein, Chp1, is required for the establishment of heterochromatin in fission yeast. *EMBO 7.* 23:3825–35
- Salgado PS, Koivunen MR, Makeyev EV, Bamford DH, Stuart DI, Grimes JM. 2006. The structure of an RNAi polymerase links RNA silencing and transcription. *PLoS Biol.* 4:e434
- Schramke V, Sheedy DM, Denli AM, Bonila C, Ekwall K, et al. 2005. RNA-interference-directed chromatin modification coupled to RNA polymerase II transcription. *Nature* 435:1275–79
- Segers GC, van Wezel R, Zhang X, Hong Y, Nuss DL. 2006. Hypovirus papain-like protease p29 suppresses RNA silencing in the natural fungal host and in a heterologous plant system. *Eukaryot. Cell* 5:896–904
- Segers GC, Zhang X, Deng F, Sun Q, Nuss DL. 2007. Evidence that RNA silencing functions as an antiviral defense mechanism in fungi. *Proc. Natl. Acad. Sci. USA* 104:12902–6
- Shanker S, Job G, George OL, Creamer KM, Shaban A, Partridge JF. 2010. Continuous requirement for the Clr4 complex but not RNAi for centromeric heterochromatin assembly in fission yeast harboring a disrupted RITS complex. *PLoS Genet.* 6:e1001174
- Shiu PK, Glass NL. 2000. Cell and nuclear recognition mechanisms mediated by mating type in filamentous ascomycetes. *Curr. Opin. Microbiol.* 3:183–88
- Shiu PK, Metzenberg RL. 2002. Meiotic silencing by unpaired DNA: properties, regulation and suppression. *Genetics* 161:1483–95
- Shiu PK, Raju NB, Zickler D, Metzenberg RL. 2001. Meiotic silencing by unpaired DNA. Cell 107:905–16

- Shiu PK, Zickler D, Raju NB, Ruprich-Robert G, Metzenberg RL. 2006. SAD-2 is required for meiotic silencing by unpaired DNA and perinuclear localization of SAD-1 RNA-directed RNA polymerase. *Proc. Natl. Acad. Sci. USA* 103:2243–48
- Son H, Min K, Lee J, Raju NB, Lee YW. 2011. Meiotic silencing in the homothallic fungus Gibberella zeae. Fungal Biol. 115:1290–302
- Sun Q, Choi GH, Nuss DL. 2009. A single Argonaute gene is required for induction of RNA silencing antiviral defense and promotes viral RNA recombination. *Proc. Natl. Acad. Sci. USA* 106:17927–32
- Suzuki N, Chen B, Nuss DL. 1999. Mapping of a hypovirus p29 protease symptom determinant domain with sequence similarity to potyvirus HC-Pro protease. *J. Virol.* 73:9478–84
- Suzuki N, Maruyama K, Moriyama M, Nuss DL. 2003. Hypovirus papain-like protease p29 functions in trans to enhance viral double-stranded RNA accumulation and vertical transmission. *J. Virol.* 77:11697– 707
- Thomson T, Lin H. 2009. The biogenesis and function of PIWI proteins and piRNAs: progress and prospect. Annu. Rev. Cell Dev. Biol. 25:355–76
- 101. Verdel A, Jia S, Gerber S, Sugiyama T, Gygi S, et al. 2004. RNAi-mediated targeting of heterochromatin by the RITS complex. *Science* 303:672–76
- Verdel A, Moazed D. 2005. RNAi-directed assembly of heterochromatin in fission yeast. FEBS Lett. 579:5872–78
- 103. Volpe T, Martienssen RA. 2011. RNA interference and heterochromatin assembly. *Cold Spring Harb. Perspect. Biol.* 3:a003731
- 104. Volpe T, Schramke V, Hamilton GL, White SA, Teng G, et al. 2003. RNA interference is required for normal centromere function in fission yeast. *Chromosome Res.* 11:137–46
- 105. Volpe TA, Kidner C, Hall IM, Teng G, Grewal SIS, Martienssen RA. 2002. Regulation of heterochromatic silencing and histone H3 lysine-9 methylation by RNAi. *Science* 297:1833–37
- 106. Wang X, Hsueh Y-P, Li W, Floyd A, Skalsky R, Heitman J. 2010. Sex-induced silencing defends the genome of *Cryptococcus neoformans* via RNAi. *Genes Dev.* 24:2566–82
- White SA, Allshire RC. 2008. RNAi-mediated chromatin silencing in fission yeast. Curr. Top. Microbiol. Immunol. 320:157–83
- 108. Wierzbicki AT, Haag JR, Pikaard CS. 2008. Noncoding transcription by RNA polymerase Pol IVb/Pol V mediates transcriptional silencing of overlapping and adjacent genes. *Cell* 135:635–48
- 109. Xiao H, Alexander WG, Hammond TM, Boone EC, Perdue TD, et al. 2010. QIP, a protein that converts duplex siRNA into single strands, is required for meiotic silencing by unpaired DNA. *Genetics* 186:119–26
- Zaratiegui M, Castel SE, Irvine DV, Kloc A, Ren J, et al. 2011. RNAi promotes heterochromatic silencing through replication-coupled release of RNA Pol II. *Nature* 479:135–38
- 111. Zhang K, Mosch K, Fischle W, Grewal SIS. 2008. Roles of the Clr4 methyltransferase complex in nucleation, spreading and maintenance of heterochromatin. *Nat. Struct. Mol. Biol.* 15:381–88
- 112. Zhang X, Nuss DL. 2008. A host Dicer is required for defective viral RNA production and recombinant virus vector RNA instability for a positive sense RNA virus. *Proc. Natl. Acad. Sci. USA* 105:16749–54
- 113. Zhang X, Segers GC, Sun Q, Deng F, Nuss DL. 2008. Characterization of hypovirus-derived small RNAs generated in the chestnut blight fungus by an inducible DCL-2-dependent pathway. *J. Virol.* 82:2613–19
- Djupedal I, Portoso M, Spåhr H, Bonilla C, Gustafsson CM, et al. 2005. RNA Pol II subunit Rpb7 promotes centromeric transcription and RNAi-directed chromatin silencing. *Genes Dev.* 19(19):2301–6

101. Identifies the RITS complex, which is the link between the RNAi machinery and heterochromatin assembly factors in *S. pombe.*

105. The first study to demonstrate that RNAi components are required for heterochromatin formation in *S. pombe*.

Annu. Rev. Microbiol. 2012.66:305-323. Downloaded from www.annualreviews.org by Universidade Federal do Parana on 03/05/13. For personal use only.

Annual Review of Microbiology

Volume 66, 2012

Contents

A Fortunate Journey on Uneven Grounds Agnes Ullmann
Memories of a Senior Scientist: On Passing the Fiftieth Anniversary of the Beginning of Deciphering the Genetic Code <i>Peter Lengyel</i>
Yeast ATP-Binding Cassette Transporters Conferring Multidrug Resistance <i>Rajendra Prasad and Andre Goffeau</i>
'Gestalt,' Composition and Function of the <i>Trypanosoma brucei</i> Editosome <i>H. Ulrich Göringer</i>
Physiology and Diversity of Ammonia-Oxidizing Archaea David A. Stahl and José R. de la Torre
Bacterial Persistence and Toxin-Antitoxin Loci Kenn Gerdes and Etienne Maisonneuve
Activating Transcription in Bacteria David J. Lee, Stephen D. Minchin, and Stephen J.W. Busby
Herpesvirus Transport to the Nervous System and Back Again Gregory Smith
A Virological View of Innate Immune Recognition Akiko Iwasaki
DNA Replication and Genomic Architecture in Very Large Bacteria Esther R. Angert
Large T Antigens of Polyomaviruses: Amazing Molecular Machines Ping An, Maria Teresa Sáenz Robles, and James M. Pipas
Peroxisome Assembly and Functional Diversity in Eukaryotic Microorganisms <i>Laurent Pieuchot and Gregory Jedd</i>

Microbial Population and Community Dynamics on Plant Roots and Their Feedbacks on Plant Communities <i>James D. Bever, Thomas G. Platt, and Elise R. Morton</i>
Bacterial Chemotaxis: The Early Years of Molecular Studies Gerald L. Hazelbauer
RNA Interference Pathways in Fungi: Mechanisms and Functions Shwu-Shin Chang, Zhenyu Zhang, and Yi Liu
Evolution of Two-Component Signal Transduction Systems Emily J. Capra and Michael T. Laub
The Unique Paradigm of Spirochete Motility and Chemotaxis Nyles W. Charon, Andrew Cockburn, Chunhao Li, Jun Liu, Kelly A. Miller, Michael R. Miller, Md. A. Motaleb, and Charles W. Wolgemuth 349
Vaginal Microbiome: Rethinking Health and DiseaseBing Ma, Larry J. Forney, and Jacques Ravel371
Electromicrobiology Derek R. Lovley
Origin and Diversification of Eukaryotes <i>Laura A. Katz</i>
Genomic Insights into Syntrophy: The Paradigm for Anaerobic Metabolic Cooperation <i>Jessica R. Sieber, Michael J. McInerney, and Robert P. Gunsalus</i>
Structure and Regulation of the Type VI Secretion System Julie M. Silverman, Yannick R. Brunet, Eric Cascales, and Joseph D. Mougous 453
Network News: The Replication of Kinetoplast DNA Robert E. Jensen and Paul T. Englund
Pseudomonas aeruginosa Twitching Motility: Type IV Pili in Action Lori L. Burrows
Postgenomic Approaches to Using Corynebacteria as Biocatalysts Alain A. Vertès, Masayuki Inui, and Hideaki Yukawa

Index

Cumulative Index of	of Contributing Authors,	Volumes 62-66	
---------------------	--------------------------	---------------	--

Errata

An online log of corrections to *Annual Review of Microbiology* articles may be found at http://micro.annualreviews.org/