

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/325157735>

# Dynamics of forest fires in the southwestern Amazon

Article in *Forest Ecology and Management* · September 2018

DOI: 10.1016/j.foreco.2018.04.041

CITATIONS

9

READS

901

6 authors, including:



**Sonaira Silva**

Universidade Federal do Acre

41 PUBLICATIONS 96 CITATIONS

[SEE PROFILE](#)



**Philip Fearnside**

Instituto Nacional de Pesquisas da Amazônia

612 PUBLICATIONS 20,906 CITATIONS

[SEE PROFILE](#)



**PAULO MAURICIO Lima De Alencastro Graça**

Instituto Nacional de Pesquisas da Amazônia

102 PUBLICATIONS 1,213 CITATIONS

[SEE PROFILE](#)



**Foster Brown**

Woods Hole Research Center

97 PUBLICATIONS 2,670 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Capacity Building for Participatory Monitoring of Changing Forests in Sustainable Use Areas of the Southwestern Brazilian Amazon [View project](#)



Inventário de Emissões Antrópicas e Sumidouros de Gases de Efeito Estufa do Estado do Acre [View project](#)

**The text that follows is a PREPRINT.**

**O texto que segue é um PREPRINT.**

Please cite as:

Favor citar como:

**da Silva, Sonaira Souza; Philip Martin  
Fearnside; Paulo Mauricio Lima de  
Alencastro Graça; Irving Foster Brown;  
Ane Alencar; Antonio Willian Flores de  
Melo. 2018. Dynamics of forest fires in the  
southwestern Amazon. *Forest Ecology  
and Management* 424: 312–322.  
<https://doi.org/10.1016/j.foreco.2018.04.041>**

ISSN: 0378-1127

Copyright: Elsevier

The original publication is available at:

O trabalho original está disponível em:

<http://www.sciencedirect.com/science/journal/03781127>

<http://www.elsevier.com.nl>

<https://doi.org/10.1016/j.foreco.2018.04.041>

1 19/04/18

2 **Dynamics of forest fires in the southwestern Amazon**

3  
4 Sonaira Souza da Silva<sup>1,2,\*</sup>; Philip Martin Fearnside<sup>2</sup>; Paulo Mauricio Lima de Alencastro  
5 Graça<sup>2</sup>; Irving Foster Brown<sup>3</sup>; Ane Alencar<sup>4</sup>; Antonio Willian Flores de Melo<sup>1,2</sup>

6  
7 <sup>1</sup> Federal University of Acre (UFAC), CEP 69.980-000, Cruzeiro do Sul, Acre, Brazil.  
8 [sonairasilva@gmail.com](mailto:sonairasilva@gmail.com); [willianflores@ufac.br](mailto:willianflores@ufac.br)

9 <sup>2</sup> National Institute for Research in Amazonia (INPA), Caixa Postal 2223, CEP 69.080-  
10 971, Manaus, Amazonas, Brazil. [philip.fearnside@gmail.com](mailto:philip.fearnside@gmail.com); [pmlag@inpa.gov.br](mailto:pmlag@inpa.gov.br)

11 <sup>3</sup> Woods Hole Research Center, 149 Woods Hole Road, Falmouth, MA 02540-1644,  
12 USA.

13 <sup>4</sup> Environmental Research Institute of Amazonia (IPAM), SHIN CA 5, Bloco J2, Sala  
14 309, CEP 71.503-505, Brasília, DF, Brazil. [ane@ipam.org.br](mailto:ane@ipam.org.br)

15  
16  
17 \*Corresponding Author: [sonairasilva@gmail.com](mailto:sonairasilva@gmail.com)

18 Postal address: UFAC, CEP 69.980-000, Cruzeiro do Sul, Acre, Brazil.

19 .

**20 Abstract**

21 The synergism between climatic change and human action has provided conditions for the  
22 occurrence of forest fires in the Amazon. We used annual mapping to reconstruct the history  
23 of fire in Brazil's state of Acre to understand the forest-fire regime over a period of 33 years  
24 (1984 to 2016). The burn-scar index (BSI) derived from the fractions of soil and of  
25 photosynthetic and non-photosynthetic material was generated by CLASlite© software using  
26 Landsat-TM and OLI satellite images. The area of forest-fire scars totaled 525,130 ha in the  
27 period analyzed. This total includes forests that fire affected only once (388,350 ha), twice  
28 (59,800 ha) and three times (5727 ha). The years 2005 and 2010 represent 90% of the total  
29 area of forest fires in Acre, coinciding with severe droughts caused by the anomalous  
30 warming of the tropical North Atlantic Ocean. The most heavily impacted portion of Acre  
31 was in the eastern part of the state, which has the greatest forest fragmentation, consolidation  
32 of agricultural activity and presence of settlement projects. In 2005, the municipalities of  
33 Acrelândia, Plácido de Castro and Senador Guiomard accounted for more than 50% of the  
34 forest remnants impacted by fire. Of the total extent of forest fires in Acre, 43% occurred in  
35 settlement projects administered by the National Institute for Colonization and Agrarian  
36 Reform (INCRA) and 16% in conservation units administered by the Ministry of  
37 Environment (MMA). The area of forest fires was 36 times greater in the 16 years after 2000,  
38 compared to the 16 years before 2000. The frequency of fires increased dramatically from one  
39 fire episode roughly every ten years (period from 1984 to 2004), to one fire every five years  
40 (period from 2005 to 2016). With the projections of warmer climate and advancing  
41 deforestation, the dynamics of forest fires in Acre will tend to be more intense and frequent.

**42 Keywords**

43 *droughts, forest degradation, fragmentation, climate change, tropical forest, rain forest,*  
44 *Brazil, Amazonia, Acre*

## 45 1. Introduction

46 Forest fires and logging have stood out as the principal causes of forest degradation in  
47 the Amazon in recent years (Aragão and Shimabukuro, 2010; Barlow et al., 2016; Bowman et  
48 al., 2009; Cochrane and Barber, 2009; Trumbore et al., 2015). Souza et al. (2013) estimated  
49 that forest degradation represents 30% of the deforested area in Brazilian Amazonia as a  
50 whole.

51 The normal conditions of the Amazonian climate, with high humidity and rainfall, do  
52 not favor the occurrence of natural fires (Fernandes et al., 2011). The Amazon has  
53 experienced Mega El Niño events accompanied by large fires at intervals ranging from 300 to  
54 500 years from 1800 to 400 BP (Bush et al., 2008; Meggers, 1994). However, in recent years  
55 the synergism between climatic extremes and human action has provided conditions for the  
56 occurrence of large forest fires at much shorter intervals than in the past (Aragão and  
57 Shimabukuro, 2010; Cochrane and Barber, 2009; Lewis et al., 2011). Over the past 40 years,  
58 forest-fire peaks have occurred every 4 to 5 years in different parts of the basin (Alencar et al.,  
59 2004; Aragão et al., 2018; Silva et al., 2013; Vasconcelos et al., 2013; Xaud et al., 2013).

60 Vulnerability of Amazonian forests to fires, as well as the spatial and temporal  
61 distributions of the fires, have been associated with extreme droughts caused by anomalous  
62 increases in sea-surface temperature in the Pacific in the case of El Niño and in the tropical  
63 North Atlantic in the case of the Atlantic dipole. The northern and northeastern portions of  
64 Brazilian Amazonia are most affected by El Niño (Alencar et al., 2004; Chen et al., 2013;  
65 Fearnside, 1990; Schroeder et al., 2009). In the southern and southeastern portions of  
66 Amazonia, extreme drought is often associated with warming the surface of the tropical  
67 portion of the North Atlantic Ocean (Aragão et al., 2018, 2007; Chen et al., 2013; Marengo et  
68 al., 2008; Zeng et al., 2008). There is also a combination of these two types of anomalous  
69 warming that has an impact throughout the Amazon, as in 1982-83, 1997-98, 2009-10 and  
70 2015/2016 (Aragão et al., 2018; Marengo and Espinoza, 2016).

71 Mapping forest-fire scars in the Amazon allows integration of temporal resolution  
72 with the geographical context of the fire. Alencar et al. (2011) mapped such scars over a 23  
73 year period at a fine spatial resolution (30-m pixels). Morton et al. (2013) monitored forest  
74 fires over a period of 13 years in southern Amazonia, but with low spatial-resolution images  
75 (250-m pixels). Mapping using political boundaries of states, which represents the level at  
76 which decisions to control and prevent deforestation and fires are taken, has been done in a  
77 few studies for specific years: Acre in 2005 (Shimabukuro et al., 2009), Mato Grosso in 2010  
78 (Anderson et al., 2015) and Roraima in 1997/98 (Barbosa and Fearnside, 1999; Barni et al.,  
79 2015). With an increasing number of mappings of forest fires, there has been an evolution in  
80 the validation and calculation of uncertainty (Anderson et al., 2017; Foody, 2008; Mack et al.,  
81 2014; Padilla et al., 2014).

82 Acre was the epicenter of the severe droughts of 2005 and 2010 (Lewis et al., 2011)  
83 that provided conditions for large forest fires. However, no historical analysis has been done  
84 with mapping of areas impacted by forest fires, their frequency and recurrence, and the  
85 relationship with human activity and extreme droughts. The objective of this study was to  
86 analyze the extent and characterize the spatial and temporal dynamics of forest fires in the  
87 state of Acre in the southwestern portion of the Brazilian Amazon. Our study was intended to  
88 answer the following questions: (1) What was the extent of forest fires over a period of 33  
89 years? (2) What extreme drought type had the greatest influence on the occurrence of forest  
90 fires, years of El Niño or of warm surface water in the tropical North Atlantic? (3) What is the  
91 relationship between deforestation and forest fires? (4) What is the spatial distribution of fires  
92 between municipalities and land-tenure types?

93

94 **2. Data and methods**95 **2.1. Study area**

96 The analysis covers the entire state of Acre (Figure 1). Acre is at the western end of  
 97 the "arc of deforestation," an area that extends from Paragominas, Pará to Rio Branco, Acre;  
 98 this area is connected by highways and is the location of most deforestation in the Brazilian  
 99 Amazon (Fearnside, 2005). In Acre, the agricultural frontier is expanding between the  
 100 municipalities of Rio Branco and Cruzeiro do Sul along Highway BR-364 and in the northern  
 101 portions of the municipalities of Porto Acre and Sena Madureira, which border on the state of  
 102 Amazonas.

103

104

105

106

107

108

109

110

111

112

113

114

115

116


117

118

119

120

121



122 Figure 1. Location of the study area, Acre state, Brazil.

123

124 **2.2. Mapping of forest fires based on Landsat satellite data**

125 Forest fires were defined in this study as those in which the crowns of the trees were  
 126 directly or indirectly affected by fire to the point that they cause a detectable impact on the  
 127 optical satellite images, representing the scar left by the fire. Many fires were probably not  
 128 detected because they were not strong enough to reach the canopy, affecting only the  
 129 understory of the forest.

130

131 To identify forest-fire scars in the state of Acre we used Landsat 5 TM (Thematic  
 132 Mapper), Landsat 7 ETM+ (Enhanced Thematic Mapper Plus) and Landsat 8 OLI  
 133 (Operational Land Imager) satellite images from 1984 to 2016. The images were accessed  
 without cost on the United States Geological Survey (USGS) website

134 (<http://earthexplorer.usgs.gov/>). The dates of the images used for processing are from  
 135 September to December. Images were used from March to June for years with cloud problems  
 136 or with very large fires in order to map the fire events in their totality (Supplementary  
 137 Material, Table S1 and Figure S1).

138 Image processing was performed using CLASlite 3.0 free software, a compact version  
 139 of the Carnegie Landsat Analysis System (CLAS). This software uses a spectral-mixing  
 140 model associated with a robust spectral library to generate fractions that represent the main  
 141 biophysical components of the landscape in a pixel (Asner et al., 2009). The gross images in  
 142 DN (digital number) were automatically corrected radiometrically for atmospheric effects and  
 143 transformed into reflectance (Figure 2a).

144

145

146

147

148

149

150

151

152


153

154

155

156

157



158 Figure 2. Example of (a) reflectance, (b) fractions and (c) Burn-scar index (BSI) in a Landsat  
 159 TM5 image from 13 October 2005, Path 002, Row 067.

160 Fraction-images were derived from the linear spectral mixing model. The following  
 161 fractions were identified: photosynthetically active vegetation (PAV), non-photosynthetic  
 162 vegetation (NPV) and soil (S). Based on the fraction images (Figure 2b), the burn-scar index  
 163 (BSI) (Figure 2c) was applied following Alencar (2010), as in Equation (1).

$$164 \text{ BSI} = (\text{PAV} - \text{NPV}) / (\text{PAV} + \text{NPV}) \quad (1)$$

165 In order to analyze only forested areas and reduce errors of commission, a  
 166 deforestation mask was applied year-by-year to the images used to calculate the BSI. This  
 167 reduces the interpositions of the NPV fraction between deforested areas and forest-fire areas,  
 168 mainly where the fire was intense.

169 For the years 1997 to 2016 a deforestation mask was applied with data from the  
 170 PRODES program of INPE (National Institute for Space Research), which monitors Brazil's  
 171 Amazon rainforest annually (Brazil, INPE, 2016). For the period from 1984 to 1986 the  
 172 image interpretation for deforestation was done by manual editing. Local knowledge of an  
 173 interpreter is necessary in order to avoid commission errors with specific punctual events,  
 174 such as blowdowns, vegetation under extreme water stress and senescent bamboo populations



175 in open forests (Carvalho et al., 2013; Espírito-Santo et al., 2014; Nelson et al., 1994)  
 176 (Supplementary Material, Figure S2).

177 We obtained the final product for forest-fire mapping by the BSI image-slicing  
 178 method, subjectively defining the thresholds by trial and error using the color composition of  
 179 the reflectance images as a guide to identify forest-fire scars. For each scene we set an  
 180 average of 10 thresholds, after which it was possible to standardize the threshold value for the  
 181 scene. Silva et al. (2013) observed that there is no standard or fixed threshold for this  
 182 identification, which changes from scene to scene according to fire intensity, vegetation  
 183 contrast, and image noise. In this study, the average threshold in all years and scenes  
 184 identified by slicing ranged from -0.15 to 0.4. In years of extreme drought the threshold  
 185 ranged from -0.3 to 0.53, while in normal years the threshold ranged from -0.1 to 0.38.

186 After slicing the BSI image, a medium smoothing filter with a window of  $5 \times 5$  pixels  
 187 was applied to reduce the number of isolated badly classified pixels. The next step was to  
 188 convert the raster map into a vector map for visual auditing, allowing deletion of poorly  
 189 sorted polygons.


190

### 191 2.3. Validation of the mapping of forest fires

192 Remote sensing has been widely used for assessing the dynamics of land use, but field  
 193 validation is indispensable in order to compare remotely sensed data with ground truth and  
 194 estimate the margin of error. Validation of mapping in a large area like the state of Acre  
 195 requires visiting a sufficient number of points in the field – a task that requires substantial  
 196 financial resources. To supplement our field points we therefore used additional points at  
 197 randomly selected locations on the images, and the classifications of these points were  
 198 confirmed by experienced interpreters.

199 Validation in the field was performed for 139 points, and an additional 2500 random  
 200 points were identified on the Landsat images (Figure 3). Verification in the field was  
 201 performed between 2015 and 2016; 22 points were checked in intact forest areas and 106  
 202 forest points were checked at locations mapped as forest-fire scars. In addition, 21 points were  
 203 observed by Irving Foster Brown in overflights in 2005. Field observations determined  
 204 whether charcoal was present at the base of standing or fallen trees, and residents confirmed  
 205 the impossibility of entering the forest. To validate the mapping using random points we used  
 206 the Landsat image base to map the 2005 scars, generating 500 points in areas mapped as scars  
 207 and 2000 points in intact forest areas. These points were audited by two interpreters.

208



216 Figure 3. Location of (a) validation points in the field and (b) random points of forest-fire  
 217 scars.



218 Assessment of the overall accuracy of the classification and estimation of errors of  
 219 omission and commission were performed using an error matrix as proposed by Anderson et  
 220 al. (2017). The accuracy of the mapping (global accuracy) was 98%; errors of omission were  
 221 0.7%, and errors of commission were 0.6% (Table 1). The main errors of omission identified  
 222 were: forest fires occurring after the date of the image used for mapping, low-intensity fires  
 223 with little impact on the canopy and fires in small forest fragments. The errors of commission  
 224 were: shading by thin clouds and smoke, displacement of the deforestation mask and  
 225 vegetation under extreme water stress.

226 Table 1. Error matrix for mapping forest-fire scars.

Classification	Burned forest	Unburned forest	Total
Mapped as burned forest	626	9	635
Mapped as unburned forest	24	2006	2030
Total	650	2015	2665
		Confidence interval	
	Estimate	Lower bound	Upper bound
Accuracy Fire forest class user's accuracy <sup>a</sup>	99%	97%	99%
Not fire forest class user's accuracy <sup>b</sup>	99%	98%	99%
Fire forest class producer's accuracy <sup>c</sup>	63%	54%	72%
Unburned forest class producer's accuracy <sup>b</sup>	100%	100%	100%
Overall accuracy	99%	98%	99%
Area error	-1%	-2%	-1%

227 a. This combines two different data sets: field observations and images.

228 b. "User's accuracy" = Probability of a pixel as classified on the map representing the  
 229 same category on the ground.

230 c. "Producer's accuracy" = Probability of a reference pixel being correctly classified.

231

#### 232 **2.4. Temporal and spatial relations of forest fires with drought, deforestation, land-use** 233 **category and logging**

##### 234 Drought

235 As an indicator to separate normal drought years from those with extreme droughts we  
 236 used satellite rainfall data from the Tropical Rainfall Measuring Mission (TRMM) and  
 237 calculated anomalies in the maximum cumulative water deficit (MCWD) for the period from  
 238 1998 to 2016 using the method of Aragão et al. (2018). We generated the MCWD for the dry  
 239 season (June to September), a period in which fire is used in agricultural practices. To  
 240 characterize the water deficit in the dry season we applied a 100-mm month<sup>-1</sup> limit to define  
 241 the dry-season months based on Aragão et al. (2007). From the MCWD anomaly we  
 242 calculated pixel-by-pixel means and standard deviations of MCWD for each year in the 1998-  
 243 2016 period, with exception of 1998, 2005 and 2010. Values of positive or negative

244 anomalies ( $\sigma$ ) for  $1.65 \leq \sigma < 1.96$  are significant at the 90% confidence level, and for  $1.96 \leq \sigma$   
245  $< 2.58$  these values are significant at the 95% confidence level.

246 To characterize the entire period of forest-fire mapping we used data from the  
247 conventional meteorological station at Rio Branco reported in the Meteorological Database  
248 for Teaching and Research (BDMEP) of the National Institute of Meteorology (INMET) for  
249 the period from 1984 to 2016. We used the maximum number of consecutive days with  
250 rainfall less than 3 mm and the number of days with maximum daily temperature above  $35^\circ$   
251 C for the dry season period, from June to October. The choice of daily maximum temperature  
252 was defined by sensitivity tests that could or differentiate years of extreme drought from other  
253 years.

254

255 Deforestation

256 We used data from the PRODES program of INPE (National Institute for Space  
257 Research) for the period from 1997 to 2016 to evaluate the reduction of forest-fire scars by  
258 subsequent deforestation. We analyzed the percentage of deforestation of scars burned only  
259 once, burned two times and burned three times. We also analyzed the time elapsed between  
260 forest burning and subsequent deforestation.

261

262 Land-use category

263 The spatial distribution of forest fires for 2005 and 2010 was analyzed in relation to  
264 the municipal (county) boundaries and land -use categories that were maintained from 2004 to  
265 2016. Municipal boundaries were obtained from the IBGE (Brazilian Institute of Geography  
266 and Statistics) database updated to 2005, containing 22 registered municipalities (Brazil,  
267 IBGE, 2016). The data on Acre's land tenure were provided by the Acre state government  
268 from the 2nd version of the Economic Ecological Zoning of Acre, which is considered to be  
269 an official database. The following classes were analyzed: INCRA settlement projects (SPs),  
270 private properties (PPs), federal government lands (GLs), conservation units (CUs) and  
271 indigenous lands (ILs). Identification of the years of fires and the delimitation of municipal  
272 boundaries and of land-use categories were carried out to determine where there were the  
273 most fires and whether these could be related to the type of land use. This also serve to avoid  
274 misleading results that could be caused by alterations of land-use categories and political  
275 boundaries, which can occur over time.

276

277 Logging

278 We used Landsat scene 002/067 as the reference for exploratory mapping of the area  
279 with logging. The identification was made visually in the period from 1984 to 2016 using the  
280 same scenes that were used to map forest fires. We analyzed four combinations: logging that  
281 did not coincide with forest fires, logging before forest fires, logging in the same year as  
282 forest fires and logging after forest fires.

283

284 **3. Results**

285 **3.1. History of forest fires**

286 The total area of forest-fire scars mapped over the 33-year period was 525,132 ha  
287 (Figures 4a and 4b), with a 95% confidence interval of 521,369 ha - 528,426 ha calculated by

288 the method of Anderson et al. (2017). This total includes forests that were affected by fire  
 289 only once (388,351 ha), twice (59,800 ha) and three times (5727 ha). The years 2005 and  
 290 2010 accounted for 471,745 ha, or 90% of the total area affected by forest fires over the 33-  
 291 year period.

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312


313

314

315

316

317



306

307

(b)

308

309

310

311

312


313

314

315

316

317



318

319

320

321

322

323

Figure 4. Historical series of forest fires for the state of Acre for the period from 1984 to 2016. (a) Spatial distribution of fires and (b) Temporal distribution of the annual area of occurrence of forest fires, where the cross-hatched bars indicate years of extreme drought from El Niño and/or anomalous warming of the tropical North Atlantic Ocean (Atlantic dipole).

324 Over the 33-year period there was an average of one large fire every seven years  
 325 (1987, 1998, 2005, 2010 and 2016). With the shortening of the recurrence time of forest fires,  
 326 the area of forest impacted by fire increased over time. The burn-scar area in the second half  
 327 of the study period (2001 to 2016) was 36 times greater than the area burned in the first half  
 328 (1984 to 2000) (Figure 4b). More detailed information on the total, maximum and average  
 329 area of forest fire scars is provided in the Supplementary Material (Table S2).

330

### 331 3.2. Extreme drought

332 The years with extreme droughts recorded by both the MCWD anomaly analysis  
 333 (Figure 5; Annual maps are provided in the Supplementary Material, Figure S3) and the  
 334 maximum number of consecutive days without rainfall and with high temperatures (Figure 6)  
 335 were: 1987, 1998, 2005, 2010 and 2016, which concentrated 98% of the forest fire scars.  
 336 These years coincide with the years with areas of forest-fire-scars greater than 6000 ha. The  
 337 MCWD anomaly in all extreme-drought years was greater than -1.96, which is compatible  
 338 with periods longer than 43 consecutive days without rainfall and 35 days with temperatures >  
 339 35 °C. The droughts of 2005 and 2010 were stronger, with a further 62 consecutive days  
 340 without rainfall, reflecting anomaly values below -3. Unlike 1987 and 1998, when forest fires  
 341 affected less than 6000 ha and were restricted to the eastern region of Acre, the El Niño of  
 342 2016 impacted four times more area and affected parts of Acre that had never been impacted  
 343 since the beginning of our monitoring, as in the municipality of Feijó in central Acre. In 2016  
 344 there was a record number of days with high temperatures (66 days), which may have  
 345 aggravated the effect of the rainfall deficit.

346

347

348

349

350

351

352

353

354

355

356

357

358

359


360

361

362

363

364



360 Figure 5. Boxplots of MCWD (maximum cumulative water deficit) for the state of Acre from  
 361 1998 to 2016.





Figure 6. Maximum numbers of consecutive days without rain each year and total number of days with temperature above 35 °C in the months from July to October, based on data from the conventional meteorological station in Rio Branco from 1984 to 2016.

Analyzing the sequence of Landsat images for 2005, 2010 and 2016 every 15 days for scenes 001/067 and 002/067, we observed that the cumulative daily water deficit (CWD) for the first visualization of forest-fire scars was from -174 mm to -200 mm in the most-fragmented area (scene 001/067). The CWD ranged from -220 mm to -230 mm in the least-fragmented area (scene 002/067). For forest-fire scars in large non-fragmented forest blocks, the CWD was -319 mm. This information can be of key importance as a basis for issuing alerts.

### 3.3. Fire and deforestation

There was no relation in the time series between fires and deforestation in the state of Acre (Figure 7). However, 88% of the forest-fires scar area was in the eastern portion of Acre, where there is more deforestation and greater forest fragmentation (Figure 5a). Another interaction observed was that 27% of the area affected by forest fires was deforested in the years following the occurrence of the fire, leaving 349,760 ha of forests in Acre still standing in 2015 that had been degraded by fire (Table 2). The smaller the area of a forest fragment affected by fire, the greater the chance of this area being subsequently. In years of normal drought, fires affect small areas as compared to extreme-drought years; however, the percentage of the area of the scars that is subsequently deforested is higher (average 68%) for scars formed in years without extreme drought than in other years (average of 30%). The time course for this deforestation also changes, and in the period from 1984 to 2001 the time for deforestation of at least 60% of the forest-fire scars area averaged 14 years, and for the subsequent period (2002-2015) the time to reach this percentage averaged six years. Up to 2016 the percentages of the burned area that had been deforested were 44% for polygons up to 50 ha in area, 26% for polygons 50 to 1000 ha in area and 16% for polygons greater than 1000 ha in area.



417 Figure 7. Annual rates of deforestation from PRODES (INPE's Amazon Monitoring Project)  
418 and area of annual forest fires in the state of Acre. Note that the "year" for PRODES is from 1  
419 August of the previous year to 31 July of the reported year, while the "year" for forest fires is  
420 from 1 January to 31 December.

421

422 Table 2. Forest area burned one, two or three times and its subsequent conversion to  
423 deforestation in the state of Acre.

Number of times burned	Area burned (ha)	% of total area burned	Maximum polygon area of burned forest (ha)	Burned forest area subsequently deforested (ha)	% Deforested of burned forest area
1	388,350	86	14,840	105,900	27
2	59,800	13	3,230	15,600	26
3	5,730	1	490	1,380	24
Total	453,880	100		104,120	27


424

### 425 3.4. Location of forest fires by land-tenure categories and geopolitical boundaries

426 Renovation and management of pastures with the use of fire were the activities found  
427 at deforested locations near 95% of the forests points visited in the process of  
428 validating the mapping. Pasture burning is the main source of ignition for fires that escaped in  
429 2005 and 2010. The type of agricultural activity is closely related to land-tenure categories. In  
430 2005 and 2010, settlement projects had the highest percentage of forest-fire scars (43% of the  
431 total area of scars: 156,600 ha in 2005 and 50,960 ha in 2010: Figure 8). Settlement projects  
432 represented the land-tenure category with the greatest use of land for agriculture and  
433 livestock. The private property category had forest fire in 28% of the total area of fire scars  
434 (95,068 ha in 2005 and 33,538 ha in 2010), conservation units had 15% (52,522 ha in 2005  
435 and 19,645 ha in 2010), federal government lands had 12% (42,626 ha and 13,960 ha) and  
436 indigenous lands has 1% (2535 ha in 2005 and 2339 ha in 2010).

437

438



439

440

441

442 Figure 8. Contribution of land-tenure categories to the area of forest fires in the state of Acre  
 443 between 1984 and 2015: (a) forest area affected by fires in five land categories and (b) forest  
 444 area remaining by 2016 affected by forest fires in each land-ownership category.

445

446 The absolute extent of the forest area affected by fire in conservation units is of  
 447 concern. In the Chico Mendes Extractive Reserve alone, fires affected an area of 50,470 ha  
 448 between 1984 and 2016, 42,700 ha of which was in 2005. Of the 19 conservation units in  
 449 Acre, nine had forest-fire scars from the 2005 fires and 12 from the 2010 fires  
 450 (Supplementary Material, Table S4). In years of extreme drought, even burning in small  
 451 agricultural areas can get out of control and turn into large forest fires. For example, in the  
 452 Chico Mendes Extractive Reserve, which is an area with little forest fragmentation, eight  
 453 polygons of burned forest larger than 1000 ha had their probable sources of ignition in  
 454 deforested areas averaging 30 ha in size.

455 The vulnerability of the landscape to forest fires is most evident when the analysis  
 456 covers the total area of forest remnants affected by fire within the municipal boundaries. The  
 457 municipalities of Plácido de Castro and Senador Guiomard had more than 50% of their area of  
 458 forest remnants burned in 2005 (Supplementary Material, Table S3). Of the total area affected  
 459 by forest fires in the 1984-2015 period, Rio Branco (26%), Senador Guiomard (11%), Sena  
 460 Madureira (10%), Xapuri (10%), Placido de Castro (9%) and Acrelândia (9%) had the  
 461 greatest areas of forest impacted by fire (Supplementary Material, Table S3). In both analyses,  
 462 all of the municipalities are in the portion of the state with the most fragmented forests  
 463 (Figure 9).

464

465

466


467

468

469

470





488 Figure 9. Forest-fire scars of areas burned one, two and three or more times in eastern Acre,  
489 Brazil between 1984 and 2016.

490

### 491 3.5. Logging

492 Using both systematic and visual methods, we identified 181 areas (polygons) with  
493 logging between 1984 and 2016. Of these, 45% coincided with areas of forest-fire scars and  
494 55% had no spatial coincidence. Among the coincident areas, 56% had logging was done  
495 before the fires (about 8 years earlier), 37% were logged after the occurrence of the fires (on  
496 average 4 years later) and only 7% were made in the same year of fire.

497

### 498 4. Discussion

499 The history of forest fires in Acre between 1984 and 2016 shows an almost binary  
500 behavior: in the normal climate years there is little occurrence of forest fires, while in the  
501 years of extreme drought several thousands of hectares of forest areas are burned. This binary  
502 behavior of forest fires demonstrates the importance of extreme droughts as a key factor  
503 driving the occurrence of fires. The years with the greatest area of forest-fire scars are  
504 associated with extreme droughts related to the anomalous surface warming of the tropical  
505 North Atlantic, especially in 2005 and 2010 (Lewis et al., 2011; Zeng et al., 2008).

506 Mapping the extent of forest fires can be affected by the spatial resolution of the  
507 satellite image used. For 2005, Shimabukuro et al. (2009) estimated 2800 km<sup>2</sup> of forest-fire  
508 scars in Acre using MODIS images, a difference of 700 km<sup>2</sup> in relation to the present study  
509 (3513 km<sup>2</sup> of forest-fire scars in 2005). Mapping with the MODIS satellite does not allow  
510 identification of areas smaller than 25 ha (Anderson et al. 2015). In the present study, areas

511 smaller than 30 ha in 2005 represented 3470 polygons, adding an area of 253 km<sup>2</sup> (7% of the  
512 total area mapped for that year).

513 In absolute terms, the impact of fire in years of extreme drought was higher in other  
514 Amazon states than it was in Acre. During the El Niño of 1997-1998 in Roraima, 11,394-  
515 13,928 km<sup>2</sup> of forest was impacted by fires (Barbosa and Fearnside, 1999). In Mato Grosso in  
516 2010 12,776-13,011 km<sup>2</sup> was impacted by fires caused mainly by the Atlantic dipole  
517 (Anderson et al., 2015). In Roraima, forest fires in 1997-1998 accounted for 7% of the state's  
518 forest area, followed by Mato Grosso in 2010 with 4% and Acre in 2005 with 3%.

519 Toomey et al. (2011) reported daytime warm anomalies of 3 °C in 2005 and 2010 in  
520 Acre, which, associated with water deficiency, explain the reduction of forest biomass. This  
521 association highlights the catastrophic effect of climate change on the dynamics of forest fires  
522 in Acre. In 2016, we appear to have experienced the limit of the forest's tolerance to drought  
523 when a large number of days with high temperatures associated with a period of 40  
524 consecutive days without rainfall was enough to make the forest flammable and impact  
525 29,455 ha of forests in the state. In periods of severe droughts trees lose more leaves, thus  
526 accumulating more litter. This, together with high temperatures and an open canopy due to the  
527 greater number of trees without leaves, changes the microclimate in the forest. Direct sunlight  
528 enters the forest interior, drying out the litter and making it highly combustible (Nepstad et  
529 al., 1999a).

530 The recurrence of forest fires accelerates forest degradation, especially when  
531 combined with fragmentation and logging, which cause drastic depletion of biodiversity and  
532 carbon stock (Barlow et al., 2016; Laurance et al., 2018). In the Acre River Valley, the  
533 recurrence of fire was greater than in other parts of the state, with forests burning as many as  
534 three times. This is a feature of fires throughout the arc of deforestation, with forests burning  
535 as many as five times, as observed in the state of Mato Grosso (Morton et al., 2013).

536 There was no relationship between the interannual variability of forest fires and  
537 deforestation, unlike the relationship observed by Barni et al. (2015) and Morton et al. (2013)  
538 (Figure 7). However the results presented differ from Morton et al. (2013) in relation to the  
539 conversion of burned forest to deforestation in the years following the fire, with a conversion  
540 percentage of 27% for Acre and an average of 8% for Southern Amazonia. This difference  
541 may be due to the size of the forest fire polygons mapped, Morton et al. (2013) mapping areas  
542 larger than 25 ha and our study in Acre mapping areas mapping areas larger than 1 ha. Van  
543 Marle et al. (2017) found that in part of the arc of deforestation only 31% of the fire events  
544 were explained by deforestation. Aragão and Shimabukuro (2010) showed that the growth of  
545 the number of ignition sources in the Amazon followed the inverse of the deforestation trend  
546 during the period of falling deforestation rates that occurred between 2004 and the time of  
547 their study, with serious implications for goals for reducing greenhouse-gas emissions.

548 In Acre, however, the portion of the state with the greatest extent of forest fires was  
549 also the area with the greatest deforestation. This deforestation left the remaining forests more  
550 vulnerable to fire entry due to fragmentation and due to more extensive forest edges. Forest  
551 fires are concentrated in the Acre River Valley, accounting for 88% of all forest areas affected  
552 by fire in the state of Acre. This is the region with the greatest consolidation of agricultural  
553 activities and a high level of forest fragmentation, and it has the main terrestrial connections  
554 to other parts of Brazil and to Peru and Bolivia: Highways BR-364 and BR-317.

555 Protected areas for forest conservation are affected by fire in the Acre River Valley.  
556 Fire scars in protected areas accounted for 14% of the entire forest-fire area between 1984 and  
557 2016. The Chico Mendes Extractive Reserve was especially affected, with forest fires in

558 isolated areas with continuous-forest polygons greater than 1000 ha in area impacted by fire,  
 559 especially in 2005 (Figure 10). Vasconcelos et al. (2013) also reported the occurrence of  
 560 forest affected by fire in indigenous lands in the southern Amazon. Human pressure from  
 561 settlement projects and roads around protected areas increases the likelihood of uncontrolled  
 562 fire reaching large areas of forest (Fearnside, 2009).

563

564

565

566

567

568

569

570

571


572

573

574

575

576



577 Figure 10. Example of the fires of 2005, size of polygons of forest fires in conservation units  
 578 under strong anthropic pressure, eastern Acre state, Brazil.

579

580

581 Logging is well known as a factor that increases the vulnerability of Amazonian  
 582 forests to subsequent fires (e.g., Nepstad et al., 1999b). Here we show that the inverse  
 583 relationship can also apply: forest fire can be associated with subsequent logging. These  
 584 results show the importance of improving and expanding the mapping of the logging in the  
 whole state of Acre and elsewhere in Amazonia.

585

586 Forest fires will likely be more frequent and more intense with further lengthening of  
 587 the dry season, which is already occurring in southwestern Amazonia (Fu et al., 2013). This  
 588 effect, which is linked to global climate change, would be worsened by the effect of continued  
 589 deforestation in further reducing rainfall in the dry season (Spracklen and Garcia-Carreras,  
 2015). These effects can be expected to cause an even more drastic impoverishment of forest  
 590 carbon stocks and biodiversity, as well as compromising policies for rewarding ecosystem  
 591 services.

592

593 With the advancement of mapping fire impact based on field data, it will be possible to  
 594 better understand the impact from climate change on forest degradation by fire, adaptation of  
 595 forest to new drought and/or fire regimes and of recovery of forest-carbon stocks (Berenguer  
 596 et al., 2014; Bustamante et al., 2016). By reducing uncertainties through monitoring with  
 597 remote sensing, it will be possible to improve carbon-cycle models and estimates of impacts  
 598 on biodiversity and climate, supporting social, economic and political decision-making  
 (Bustamante et al., 2016; Olofsson et al., 2013).

599 Maintaining the goals of reducing deforestation does not guarantee the maintenance of  
600 forest ecosystem services, since forest degraded by logging and fire retains only 46-61% of its  
601 potential conservation value (Barlow et al., 2016). We observed two situations combining  
602 logging and forest fires, pre-fire exploration and post-fire exploration. The first is more  
603 common in most of the Amazon (Barlow et al., 2006; Berenguer et al., 2014). In Acre the  
604 most common situation is the second. Through rapid assessment of Landsat-TM scene  
605 002/067, 26% of the identified logging occurred up to three years after the fires of 2005 or  
606 2010, unleashing a new mode of forest degradation that could further hamper forest recovery  
607 after the fires.

608 Public policies for reducing greenhouse-gas emissions have so far only considered the  
609 impact of deforestation, s in the Acre state government's program to reduce greenhouse-gas  
610 emissions (Neves et al., 2013). We recommend that future studies include calculation of  
611 emissions from forest fires over a historical series, as these fires can be a significant emissions  
612 source, especially in drought years.

613

## 614 **Conclusions**

615 Monitoring of forest fires in the state of Acre over a 33-year period (1984-2016)  
616 shows that higher frequencies and area of forest-fire scars are concentrated in the last 12  
617 years, when 95% of the area burned over the entire period was in the years 2005, 2010 and  
618 2016.

619 Forest fires are responses to drought events, increased fragmentation, and the use of  
620 fire in agricultural practices. In the eastern portion of Acre, the municipalities of Acrelândia  
621 and Senador Guimard had more than 50% of their remaining forests affected by fire in 2005.

622 The impact of humans on initiating forest fires is decisive: 42% of forest fires occurred  
623 in INCRA settlement projects and 87% of the fires occurred in the eastern portion of the state,  
624 which is the most densely populated region and accounts for 65% of deforestation in Acre.  
625 Conservation units accounted for 14% of the entire forest-fire area in Acre, showing the  
626 fragility of the landscape even with low deforestation pressure.

627 The implications of forest fires for policies to reduce carbon emissions and pay for  
628 environmental services are enormous. The area affected by forest fires in Acre in the period  
629 from 2004 to 2015 is equal to the entire area deforested in the state over the same period.

630 The joint analysis of timber extraction on increasing susceptibility of the forest to fire  
631 and of predicted increases in climatic extremes will be important to understanding the future  
632 evolution of forest fires in southwestern Amazonia.

633

## 634 **Acknowledgments**

635 We received financing from the Acre State Research Support Foundation (Call 03/2013), the  
636 National Institute of Science and Technology of Environmental Services of the Amazon  
637 (INCT-SERVAMB) (CNPq 610042/2009-2) and the Coordination for Improvement of Higher  
638 Education Personnel (CAPES) through the INPA/UFAC Interinstitutional Doctoral Program  
639 (No. 459/2013).

640

641 **References**

- 642 Alencar, A., 2010. Spatial and temporal determinants of forest fires on the Amazonian  
643 deforestation frontier: Implications for current and future carbon emissions (Ph.D.).  
644 University of Florida, Gainesville, Florida. <http://ufdc.ufl.edu/UFE0042496/00001>
- 645 Alencar, A., Asner, G.P., Knapp, D., Zarin, D., 2011. Temporal variability of forest fires in  
646 eastern Amazonia. *Ecol. Appl.* 21, 2397–2412. <https://doi.org/10.1890/10-1168.1>
- 647 Alencar, A.A.C., Solórzano, L.A., Nepstad, D.C., 2004. Modeling forest understory fires in an  
648 eastern Amazonian landscape. *Ecol. Appl.* 14, 139–149. <https://doi.org/10.1890/01-6029>
- 650 Anderson, L.O., Aragão, L.E.O.C., Gloor, M., Arai, E., Adami, M., Saatchi, S.S., Malhi, Y.,  
651 Shimabukuro, Y.E., Barlow, J., Berenguer, E., Duarte, V., 2015. Disentangling the  
652 contribution of multiple land covers to fire-mediated carbon emissions in Amazonia  
653 during the 2010 drought. *Glob. Biogeochem. Cycles* 29, 1739–1753.  
654 <https://doi.org/10.1002/2014GB005008>
- 655 Anderson, L.O., Cheek, D., Aragao, L.E., Andere, L., Duarte, B., Salazar, N., Lima, A.,  
656 Duarte, V., Arai, E., 2017. Development of a point-based method for map validation  
657 and confidence interval estimation: A case study of burned areas in Amazonia. *J.*  
658 *Remote Sens. GIS* 06. <https://doi.org/10.4172/2469-4134.1000193>
- 659 Aragão, L.E.O.C., Anderson, L.O., Fonseca, M.G., Rosan, T.M., Vedovato, L.B., Wagner,  
660 F.H., Silva, C.V.J., Junior, C.H.L.S., Arai, E., Aguiar, A.P., Barlow, J., Berenguer, E.,  
661 Deeter, M.N., Domingues, L.G., Gatti, L., Gloor, M., Malhi, Y., Marengo, J.A.,  
662 Miller, J.B., Phillips, O.L., Saatchi, S., 2018. 21st Century drought-related fires  
663 counteract the decline of Amazon deforestation carbon emissions. *Nat. Commun.* 9,  
664 536. <https://doi.org/10.1038/s41467-017-02771-y>
- 665 Aragão, L.E.O.C., Malhi, Y., Roman-Cuesta, R.M., Saatchi, S., Anderson, L.O.,  
666 Shimabukuro, Y.E., 2007. Spatial patterns and fire response of recent Amazonian  
667 droughts. *Geophys. Res. Lett.* 34, 1–5. <https://doi.org/10.1029/2006GL028946>
- 668 Aragão, L.E.O.C., Poulter, B., Barlow, J.B., Anderson, L.O., Malhi, Y., Saatchi, S., Phillips,  
669 O.L., Gloor, E., 2014. Environmental change and the carbon balance of Amazonian  
670 forests. *Biol. Rev.* 1–19. <https://doi.org/10.1111/brv.12088>
- 671 Aragão, L.E.O.C., Shimabukuro, Y.E., 2010. The incidence of fire in Amazonian forests with  
672 implications for REDD. *Science* 328, 1275–1278.  
673 <https://doi.org/10.1126/science.1186925>
- 674 Asner, G.P., Knapp, D.E., Balaji, A., Paez-Acosta, G., 2009. Automated mapping of tropical  
675 deforestation and forest degradation: CLASlite. *J. Appl. Remote Sens.* 3, 1–24.  
676 <https://doi.org/10.1117/1.3223675>
- 677 Barbosa, R.I., Fearnside, P.M., 1999. Incêndios na Amazônia Brasileira: estimativa da  
678 emissão de gases do efeito estufa pela queima de diferentes ecossistemas de Roraima  
679 na passagem do evento “El Niño” (1997/98). *Acta Amaz.* 29, 513–534.  
680 <https://doi.org/10.1590/1809-43921999294534>
- 681 Barlow, J., Lennox, G.D., Ferreira, J., Berenguer, E., Lees, A.C., Nally, R.M., Thomson, J.R.,  
682 Ferraz, S.F. de B., Louzada, J., Oliveira, V.H.F., Parry, L., Ribeiro de Castro Solar, R.,  
683 Vieira, I.C.G., Aragão, L.E.O.C., Begotti, R.A., Braga, R.F., Cardoso, T.M., Jr, R.C.  
684 de O., Souza Jr, C.M., Moura, N.G., Nunes, S.S., Siqueira, J.V., Pardini, R., Silveira,  
685 J.M., Vaz-de-Mello, F.Z., Veiga, R.C.S., Venturieri, A., Gardner, T.A., 2016.

- 686 Anthropogenic disturbance in tropical forests can double biodiversity loss from  
687 deforestation. *Nature* 535, 144–147. <https://doi.org/10.1038/nature18326>
- 688 Barlow, J., Peres, C.A., Henriques, L.M.P., Stouffer, P.C., Wunderle, J.M., 2006. The  
689 responses of understorey birds to forest fragmentation, logging and wildfires: An  
690 Amazonian synthesis. *Biol. Conserv.* 128, 182–192.  
691 <https://doi.org/10.1016/j.biocon.2005.09.028>
- 692 Barni, P.E., Pereira, V.B., Manzi, A.O., Barbosa, R.I., 2015. Deforestation and forest fires in  
693 Roraima and their relationship with phytoclimatic regions in the northern Brazilian  
694 Amazon. *Environ. Manage.* 55, 1124–1138. <https://doi.org/10.1007/s00267-015-0447-7>  
695
- 696 Berenguer, E., Ferreira, J., Gardner, T.A., Aragão, L.E.O.C., de Camargo, P.B., Cerri, C.E.,  
697 Durigan, M., Oliveira, R.C.D., Vieira, I.C.G., Barlow, J., 2014. A large-scale field  
698 assessment of carbon stocks in human-modified tropical forests. *Glob. Change Biol.*  
699 20, 3713–3726. <https://doi.org/10.1111/gcb.12627>
- 700 Bowman, D.M.J.S., Balch, J.K., Artaxo, P., Bond, W.J., Carlson, J.M., Cochrane, M.A.,  
701 D’Antonio, C.M., DeFries, R.S., Doyle, J.C., Harrison, S.P., Johnston, F.H., Keeley,  
702 J.E., Krawchuk, M.A., Kull, C.A., Marston, J.B., Moritz, M.A., Prentice, I.C., Roos,  
703 C.I., Scott, A.C., Swetnam, T.W., Werf, G.R. van der, Pyne, S.J., 2009. Fire in the  
704 Earth System. *Science* 324, 481–484. <https://doi.org/10.1126/science.1163886>
- 705 Brazil, IBGE (Instituto Brasileiro de Geografia e Estatística), 2016. Municipal boundaries  
706 limits. [https://mapas.ibge.gov.br/bases-e-referenciais/bases-cartograficas/malhas-  
707 digitais.html](https://mapas.ibge.gov.br/bases-e-referenciais/bases-cartograficas/malhas-digitais.html)
- 708 Bush, M.B., Silman, M.R., McMichael, C., Saatchi, S., 2008. Fire, climate change and  
709 biodiversity in Amazonia: A late-Holocene perspective. *Philos. Trans. R. Soc. B Biol.*  
710 *Sci.* 363, 1795–1802. <https://doi.org/10.1098/rstb.2007.0014>
- 711 Bustamante, M.M.C., Roitman, I., Aide, T.M., Alencar, A., Anderson, L.O., Aragão, L.,  
712 Asner, G.P., Barlow, J., Berenguer, E., Chambers, J., Costa, M.H., Fanin, T., Ferreira,  
713 L.G., Ferreira, J., Keller, M., Magnusson, W.E., Morales-Barquero, L., Morton, D.,  
714 Ometto, J.P.H.B., Palace, M., Peres, C.A., Silvério, D., Trumbore, S., Vieira, I.C.G.,  
715 2016. Toward an integrated monitoring framework to assess the effects of tropical  
716 forest degradation and recovery on carbon stocks and biodiversity. *Glob. Change Biol.*  
717 22, 92–109. <https://doi.org/10.1111/gcb.13087>
- 718 Carvalho, A.L., Nelson, B.W., Bianchini, M.C., Plagnol, D., Kuplich, T.M., Daly, D.C., 2013.  
719 Bamboo-dominated forests of the southwest Amazon: detection, spatial extent, life  
720 cycle length and flowering waves. *PLoS One* 8, 1–13.  
721 <https://doi.org/10.1371/journal.pone.0054852>
- 722 Chen, Y., Randerson, J.T., Morton, D.C., DeFries, R.S., Collatz, G.J., Kasibhatla, P.S.,  
723 Giglio, L., Jin, Y., Marlier, M.E., 2011. Forecasting fire season severity in South  
724 America using sea surface temperature anomalies. *Science* 334, 787–791.  
725 <https://doi.org/10.1126/science.1209472>
- 726 Chen, Y., Velicogna, I., Famiglietti, J.S., Randerson, J.T., 2013. Satellite observations of  
727 terrestrial water storage provide early warning information about drought and fire  
728 season severity in the Amazon. *J. Geophys. Res.-Biogeosciences* 118, 495–504.  
729 <https://doi.org/10.1002/jgrg.20046>



- 730 Cochrane, M.A., Barber, C.P., 2009. Climate change, human land use and future fires in the  
731 Amazon. *Glob. Change Biol.* 15, 601–612. <https://doi.org/10.1111/j.1365->  
732 2486.2008.01786.x
- 733 Espírito-Santo, F.D.B., Gloor, M., Keller, M., Malhi, Y., Saatchi, S., Nelson, B., Junior,  
734 R.C.O., Pereira, C., Lloyd, J., Frolking, S., Palace, M., Shimabukuro, Y.E., Duarte, V.,  
735 Mendoza, A.M., López-González, G., Baker, T.R., Feldpausch, T.R., Brienen, R.J.W.,  
736 Asner, G.P., Boyd, D.S., Phillips, O.L., 2014. Size and frequency of natural forest  
737 disturbances and the Amazon forest carbon balance. *Nat. Commun.* 5, 3434.  
738 <https://doi.org/10.1038/ncomms4434>
- 739 Fearnside, P.M., 1990. Fire in the tropical rain forest of the Amazon basin, in: Goldammer,  
740 J.G. (Ed.), *Fire in the Tropical Biota*, Ecological Studies. Springer Berlin &  
741 Heidelberg, pp. 106–116.
- 742 Fearnside, P.M., 2005. Deforestation in Brazilian Amazonia: History, rates and consequences.  
743 *Conserv Biol* 19(3), 680–688. <https://doi.org/10.1111/j.1523-1739.2005.00697.x>
- 744 Fearnside, P.M., 2009. *A Floresta Amazônica nas mudanças globais*, 2nd ed, INPA. INPA,  
745 Manaus, Amazonas.
- 746 Fernandes, K., Baethgen, W., Bernardes, S., DeFries, R., DeWitt, D.G., Goddard, L., Lavado,  
747 W., Lee, D.E., Padoch, C., Pinedo-Vasquez, M., Uriarte, M., 2011. North Tropical  
748 Atlantic influence on western Amazon fire season variability. *Geophys. Res. Lett.* 38,  
749 L12701. <https://doi.org/10.1029/2011GL047392>
- 750 Foody, G.M., 2008. Harshness in image classification accuracy assessment. *Int. J. Remote*  
751 *Sens.* 29, 3137–3158. <https://doi.org/10.1080/01431160701442120>
- 752 Fu, R., Yin, L., Li, W., Arias, P.A., Dickinson, R.E., Huang, L., Chakraborty, S., Fernandes,  
753 K., Liebmann, B., Fisher, R., Myneni, R.B., 2013. Increased dry-season length over  
754 southern Amazonia in recent decades and its implication for future climate projection.  
755 *Proc. Natl. Acad. Sci. U. S. A.* 110, 18110–18115.  
756 <https://doi.org/10.1073/pnas.1302584110>
- 757 Kintisch, E., 2016. How a ‘Godzilla’ El Niño shook up weather forecasts. *Science* 352, 1501–  
758 1502. <https://doi.org/10.1126/science.352.6293.1501>
- 759 Laurance, W.F., Camargo, J.L.C., Fearnside, P.M., Lovejoy, T.E., Williamson, G.B.,  
760 Mesquita, R.C.G., Meyer, C.F.J., Bobrowiec, P.E.D. Laurance, S.G.W., 2018. An  
761 Amazonian rainforest and its fragments as a laboratory of global change. *Biological*  
762 *Reviews* 93(1), 223–247. <https://doi.org/10.1111/brv.12343>
- 763 Lewis, S.L., Brando, P.M., Phillips, O.L., Heijden, G.M.F. van der, Nepstad, D.C., 2011. The  
764 2010 Amazon drought. *Science* 331, 554–554.  
765 <https://doi.org/10.1126/science.1200807>
- 766 Mack, B., Roscher, R., Waske, B., 2014. Can I trust my one-class classification? *Remote*  
767 *Sens.* 6, 8779–8802. <https://doi.org/10.3390/rs6098779>
- 768 Marengo, J.A., Espinoza, J.C., 2016. Extreme seasonal droughts and floods in Amazonia:  
769 causes, trends and impacts. *Int. J. Climatol.* 36, 1033–1050.  
770 <https://doi.org/10.1002/joc.4420>
- 771 Marengo, J.A., Nobre, C.A., Tomasella, J., Oyama, M.D., Sampaio de Oliveira, G., de  
772 Oliveira, R., Camargo, H., Alves, L.M., Brown, I.F., 2008. The drought of Amazonia  
773 in 2005. *J. Clim.* 21, 495–516. <https://doi.org/10.1175/2007JCLI1600.1>



- 774 Meggers, B.J., 1994. Archeological evidence for the impact of mega-Niño events on  
775 Amazonia during the past two millennia. *Clim. Change* 28, 321–338.  
776 <https://doi.org/10.1007/BF01104077>
- 777 Morton, D.C., Page, Y.L., DeFries, R.S., Collatz, G.J., Hurtt, G.C., 2013. Understorey fire  
778 frequency and the fate of burned forests in southern Amazonia. *Philos. Trans. R. Soc.  
779 B Biol. Sci.* 368. <https://doi.org/10.1098/rstb.2012.0163>
- 780 Nelson, B.W., Kapos, V., Adams, J.B., Oliveira, W.J., Braun, O.P.G., 1994. Forest  
781 disturbance by large blowdowns in the Brazilian Amazon. *Ecology* 75, 853–858.  
782 <https://doi.org/10.2307/1941742>
- 783 Nepstad, D.C., Moreira, A.G., Alencar, A.A.C., 1999a. *Flames in the rain forest: origins,  
784 impacts and alternatives to Amazonian fire. Pilot Program to Conserve the Brazilian  
785 Rain Forest and World Bank, Brasília.* 186 pp.  
786 [http://documents.worldbank.org/curated/en/522521468013876752/pdf/635120WP0Fla  
787 me00Box0361520B0PUBLIC0.pdf](http://documents.worldbank.org/curated/en/522521468013876752/pdf/635120WP0Flame00Box0361520B0PUBLIC0.pdf)
- 788 Nepstad, D.C., Verissimo, A., Alencar, A., Nobre, C., Lima, E., Lefebvre, P., Schlesinger, P.,  
789 Potter, C., Moutinho, P., Mendoza, E., Cochrane, M., Brooks, V., 1999b. Large-scale  
790 impoverishment of Amazonian forests by logging and fire. *Nature* 398, 505–508. d  
791 <https://doi.org/10.1038/19066>
- 792 Neves, R.F., Leal, M.J. de L.R., Vaz, F., 2013. Programa de Incentivos a Serviços Ambientais  
793 do Carbono do Estado do Acre (Programa ISA Carbono do Acre). IMC, Rio Branco.  
794 [https://mer.markit.com/br-  
795 reg/services/processDocument/downloadDocumentById/10300000029314](https://mer.markit.com/br-reg/services/processDocument/downloadDocumentById/10300000029314)
- 796 Olofsson, P., Foody, G.M., Stehman, S.V., Woodcock, C.E., 2013. Making better use of  
797 accuracy data in land change studies: Estimating accuracy and area and quantifying  
798 uncertainty using stratified estimation. *Remote Sens. Environ.* 129, 122–131.  
799 <https://doi.org/10.1016/j.rse.2012.10.031>
- 800 Padilla, M., Stehman, S.V., Litago, J., Chuvieco, E., 2014. Assessing the temporal stability of  
801 the accuracy of a time series of burned area products. *Remote Sens.* 6, 2050–2068.  
802 <https://doi.org/10.3390/rs6032050>
- 803 Panisset, J.S., Libonati, R., Gouveia, C.M.P., Machado-Silva, F., França, D.A., França,  
804 J.R.A., Peres, L.F., 2017. Contrasting patterns of the extreme drought episodes of  
805 2005, 2010 and 2015 in the Amazon Basin: Extreme droughts episodes of 2005, 2010  
806 and 2015 in the Amazon. *Int. J. Climatol.* <https://doi.org/10.1002/joc.5224>
- 807 Schroeder, W., Alencar, A., Arima, E., Setzer, A., 2009. The spatial distribution and  
808 interannual variability of fire in Amazonia, in: Keller, M., Bustamante, M., Gash, J.,  
809 Dias, P.S. (Eds.), *Amazonia and Global Change.* American Geophysical Union,  
810 Washington, DC, pp. 43–60.
- 811 Shimabukuro, Y.E., Duarte, V., Arai, E., Freitas, R.M., Lima, A., Valeriano, D.M., Brown,  
812 I.F., Maldonado, M.L.R., 2009. Fraction images derived from Terra Modis data for  
813 mapping burnt areas in Brazilian Amazonia. *Int. J. Remote Sens.* 30, 1537–1546.  
814 <https://doi.org/10.1080/01431160802509058>
- 815 Silva, S.S., Alencar, A.A., Mendoza, E., Brown, I.F., 2013. Dinâmica dos incêndios florestais  
816 no Estado do Acre nas décadas de 90 e 00, in: *Simpósio Brasileiro de Sensoriamento  
817 Remoto (SBSR), 16.* INPE, São José dos Campos, pp. 8799–8806.

- 818 Souza, J., Siqueira, J.V., Sales, M.H., Fonseca, A.V., Ribeiro, J.G., Numata, I., Cochrane,  
819 M.A., Barber, C.P., Roberts, D.A., Barlow, J., 2013. Ten-Year Landsat Classification  
820 of Deforestation and Forest Degradation in the Brazilian Amazon. *Remote Sens.* 5,  
821 5493–5513. <https://doi.org/10.3390/rs5115493>
- 822 Spracklen, D.V., Garcia-Carreras, L., 2015. The impact of Amazonian deforestation on  
823 Amazon basin rainfall. *Geophys. Res. Lett.* 42, 9546–9552.  
824 <https://doi.org/10.1002/2015GL066063>
- 825 Toomey, M., Roberts, D.A., Still, C., Goulden, M.L., McFadden, J.P., 2011. Remotely sensed  
826 heat anomalies linked with Amazonian forest biomass declines. *Geophys. Res. Lett.*  
827 38, L19704. <https://doi.org/10.1029/2011GL049041>
- 828 Trumbore, S., Brando, P., Hartmann, H., 2015. Forest health and global change. *Science* 349,  
829 814–818. <https://doi.org/10.1126/science.aac6759>
- 830 van Marle, M.J.E., Field, R.D., van der Werf, G.R., Estrada de Wagt, I.A., Houghton, R.A.,  
831 Rizzo, L.V., Artaxo, P., Tsigaridis, K., 2017. Fire and deforestation dynamics in  
832 Amazonia (1973–2014). *Glob. Biogeochem. Cycles* 2016GB005445.  
833 <https://doi.org/10.1002/2016GB005445>
- 834 Vasconcelos, S.S. de, Fearnside, P.M., Graça, P.M.L. de A., Nogueira, E.M., Oliveira, L.C.  
835 de, Figueiredo, E.O., 2013. Forest fires in southwestern Brazilian Amazonia:  
836 Estimates of area and potential carbon emissions. *For. Ecol. Manag.* 291, 199–208.  
837 <https://doi.org/10.1016/j.foreco.2012.11.044>
- 838 Xaud, H.A.M., Martins, F. da S.R.V., Santos, J.R. dos, 2013. Tropical forest degradation by  
839 mega-fires in the northern Brazilian Amazon. *For. Ecol. Manag.* 294, 97–106.  
840 <https://doi.org/10.1016/j.foreco.2012.11.036>
- 841 Zeng, N., Yoon, J.-H., Marengo, J.A., Subramaniam, A., Nobre, C.A., Mariotti, A., Neelin,  
842 J.D., 2008. Causes and impacts of the 2005 Amazon drought. *Environ. Res. Lett.* 3,  
843 014002. <https://doi.org/10.1088/1748-9326/3/1/014002>
- 844

## Supplementary Material

### Dynamics of forest fires in the southwestern Amazon

Sonaira Souza da Silva<sup>1,2,\*</sup>; Philip Martin Fearnside<sup>2</sup>; Paulo Mauricio Lima de Alencastro Graça<sup>2</sup>; Irving Foster Brown<sup>3</sup>; Ane Alencar<sup>4</sup>; Antonio Willian Flores de Melo<sup>1,2</sup>

<sup>1</sup> Federal University of Acre (UFAC), Estrada do Canela Fina, km 12, CEP 69.980-000, Cruzeiro do Sul, Acre, Brazil. [sonairasilva@gmail.com](mailto:sonairasilva@gmail.com); [willianflores@ufac.br](mailto:willianflores@ufac.br)

<sup>2</sup> National Institute for Research in Amazonia (INPA), Caixa Postal 2223, CEP 69.080-971, Manaus, Amazonas, Brazil. [philip.fearnside@gmail.com](mailto:philip.fearnside@gmail.com); [pmlag@inpa.gov.br](mailto:pmlag@inpa.gov.br)

<sup>3</sup> Woods Hole Research Center (WHRC), 149 Woods Hole Road, Falmouth, MA 02540-1644, USA. [fbrown@whrc.org](mailto:fbrown@whrc.org)

<sup>4</sup> Environmental Research Institute of Amazonia (IPAM), SHIN CA 5, Bloco J2, Sala 309, CEP 71.503-505, Brasília, DF, Brazil. [ane@ipam.org.br](mailto:ane@ipam.org.br)

\*Corresponding Author: [sonairasilva@gmail.com](mailto:sonairasilva@gmail.com)

Postal address: UFAC, CEP 69.980-000, Cruzeiro do Sul, Acre, Brazil

#### Contents of this file

Table S1. Data for the Landsat images used in this study for the mapping of forest fires in the state of Acre.

Table S2. Total, maximum and average annual area of forest fire scars for the state of Acre from 1984 to 2016.

Table S3. List of protected areas with forest-fire scars in 2005 and 2010.

Table S4. Areas of forest fires by municipality in the state of Acre in the 1984 - 2016 period, classified by the number of times each pixel had been burned.

Figure S1. Temporal distribution of hot spots between June and November in the period from 2000 to 2016

Figure S2. Main types of confusion during the mapping of forest fires.

Figure S3. Annual spatial distribution of the Maximum Cumulative Water Deficit (MCWD) anomaly for the state of Acre from 1998 to 2016.

Table S1. Data for the Landsat images used in this study for the mapping of forest fires in the state of Acre

<b>Year</b>	<b>001/067</b>	<b>002/066</b>	<b>002/067</b>	<b>002/068</b>	<b>003/066</b>	<b>003/067</b>	<b>004/065</b>	<b>004/066</b>	<b>004/067</b>	<b>005/065</b>	<b>005/066</b>	<b>006/065</b>
1984	15-Dec	17-Sep	01-Sep	06-Dec	cloud	10-Oct	15-Sep	01-Oct	01-Oct	06-Sep	cloud	cloud
1985	16-Nov	04-Sep	06-Oct	06-Oct	cloud	30-Nov	cloud	cloud	cloud	09-Sep	09-Sep	cloud
1986	cloud	10-Nov	10-Nov	25-Oct	30-Sep	30-Sep	07-Oct	07-Oct	07-Oct	14-Oct	14-Oct	cloud
1987	05-Oct	12-Oct	12-Oct	12-Oct	cloud	03-Oct	26-Oct	cloud	cloud	15-Sep	01-Oct	cloud
1988	05-Set	14-Oct	01-Oct	30-Oct	cloud	cloud	10-Sep	10-Sep	cloud	cloud	cloud	26-Oct
1989	24-Sep	15-Sep	cloud	15-Sep	22-Sep	22-Sep	13-Sep	cloud	cloud	25-Dec	04-Sep	30-Nov
1990	14-Nov	cloud	18-Sep	18-Sep	cloud	cloud	cloud	16-Sep	16-Sep	cloud	07-Sep	cloud
1991	30-Sep	10-Dec	10-Dec	10-Dec	15-Oct	14-Oct	cloud	05-Oct	05-Oct	cloud	cloud	cloud
1992	02-Oct	10-Nov	23-Sep	23-Sep	30-Sep	30-Sep	30-Sep	30-Sep	05-Sep	28-Sep	28-Sep	cloud
1993	06-Nov	25-Aug	25-Aug	25-Aug	20-Nov	04-Nov	08-Sep	cloud	08-Sep	30-Aug	30-Aug	08-Oct
1994	11-Dec	29-Sep	29-Sep	cloud	cloud	cloud	14-Nov	14-Nov	14-Nov	05-Nov	cloud	25-Sep
1995	cloud	06-Sep	18-Oct	18-Oct	23-Sep	23-Sep	cloud	14-Sep	cloud	05-Sep	07-Oct	27-Aug
1996	16-Dec	cloud	cloud	20-Oct	09-Sep	09-Sep	02-Oct	02-Oct	02-Oct	26-Nov	26-Nov	14-Oct
1997	30-Sep	21-Sep	05-Sep	08-Nov	28-Sep	12-Sep	cloud	19-Sep	05-Oct	12-Oct	12-Oct	17-Sep
1998	15-Sep	10-Oct	29-Dec	24-Sep	30-Aug	30-Aug	22-Sep	22-Sep	22-Sep	13-Sep	13-Sep	04-Sep
1999	16-Sep	05-Oct	05-Oct	05-Oct	18-Sep	18-Sep	11-Oct	11-Oct	11-Oct	03-Nov	03-Nov	23-Sep
2000	16-Dec	24-Nov	18-Dec	18-Dec	04-Sep	cloud	27-Sep	25-Sep	25-Sep	04-Oct	02-Sep	11-Oct
2001	25-Sep	08-Sep	02-Oct	02-Oct	09-Oct	09-Oct	cloud	cloud	cloud	07-Oct	07-Oct	15-Nov
2002	07-Nov	26-Aug	26-Aug	26-Aug	04-Oct	04-Oct	27-Oct	11-Oct	11-Oct	16-Sep	16-Sep	23-Sep
2003	17-Oct	09-Nov	23-Oct	24-Oct	15-Oct	15-Oct	cloud	cloud	cloud	cloud	cloud	25-Feb
2004	01-Sep	08-Sep	24-Jun	26-Oct	15-Sep	17-Oct	09-Nov	09-Nov	09-Nov	31-Oct	28-Aug	23-Nov
2005	20-Sep	13-Oct	11-Sep	13-Oct	05-Nov	05-Nov	25-Sep	25-Sep	25-Sep	16-Sep	16-Sep	23-Sep
	22-Oct	10-Jun	13-Oct	25-May	16-May							
2006	07-Sep	14-Sep	14-Sep	13-Aug	21-Sep	05-Sep	28-Sep	28-Sep	28-Sep	03-Sep	03-Sep	25-Aug
2007	26-Sep	16-Aug	17-Sep	06-Sep	24-Sep	23-Aug	29-Jul	29-Jul	11-Jun	22-Sep	22-Sep	cloud
2008	28-Sep	18-Aug	18-Aug	19-Sep	28-Oct	28-Oct	01-Sep	16-Aug	01-Sep	24-Sep	24-Sep	14-Aug
2009	01-Oct	06-Sep	06-Sep	06-Sep	13-Sep	13-Sep	04-Sep	04-Sep	04-Sep	14-Nov	14-Nov	04-Oct
2010	04-Oct	11-Oct	13-Oct	10-Oct	03-Nov	03-Nov	06-Aug	07-Sep	07-Sep	30-Sep	30-Sep	20-Aug
	01-Jun	24-Jun	24-Jun	24-Jun	30-May	30-May		05-May		15-Jul		
2011	05-Sep	30-Oct	30-Oct	30-Oct	18-Aug	18-Aug	09-Aug	09-Aug	22-Aug	15-Aug	15-Aug	08-Sep
2012	02-Nov	06-Sep	06-Sep	06-Sep	13-Sep	29-Sep		20-Sep	20-Sep	11-Sep	11-Sep	18-Sep

2013	26-Sep	17-Sep	17-Sep	17-Sep	11-Nov	08-Sep	30-Aug	30-Aug	01-Oct	11-Dec	11-Dec	13-Sep
2014	29-Sep	07-Nov	22-Oct	22-Oct	13-Oct	13-Oct	18-Sep	18-Sep	02-Sep	24-Aug	25-Sep	18-Oct
2015	18-Oct	09-Oct	12-Dec	12-Dec	17-Nov	03-Dec	23-Oct	23-Oct	23-Oct	12-Sep	12-Sep	21-Oct
2016	09-Sep	09-Sep	09-Sep	09-Sep	18-Oct	15-Aug	07-Sep	07-Sep	07-Sep	29-Aug	14-Sep	16-Aug
	05-Nov	28-Nov	09-Sep	09-Sep	18-Oct	19-Nov				01-Nov	01-Nov	

Table S2. Total, maximum and average annual area of forest fire scars for the State of Acre from 1984 to 2016.

Year	Total	Maximum	Average
1984	15	8	4
1985	5	4	2
1986	51	22	10
1987	6,206	1,012	14
1988	334	56	7
1989	117	33	10
1990	642	47	12
1991	30	13	10
1992	98	51	24
1993	159	25	11
1994	295	33	7
1995	662	93	12
1996	23	10	8
1997	167	62	12
1998	5,892	215	12
1999	330	29	7
2000	178	28	7
2001	350	142	18
2002	261	70	16
2003	1,909	121	14
2004	447	48	8
2005	351,285	15,040	73
2006	1,160	164	6
2007	525	134	10
2008	1,298	146	10
2009	272	55	6
2010	120,459	4,764	39
2011	1,021	186	16
2012	295	68	11
2013	217	28	6
2014	67	15	6
2015	904	86	7
2016	29,455	3,926	39

Table S3. List of protected areas with forest fire scars in 2005 and 2010.

<b>Conservation Units</b>	<b>2005</b>	<b>2010</b>
APA Amapá	517	72
APA Irineu Serra	66	120
APA São Francisco	7,609	3,584
Arie Seringal Nova Esperança	20	34
Floresta Estadual do Antimary	176	130
Floresta Estadual do Chandles	18	91
Floresta Nacional Santa Rosa do Purus	47	409
Parque Nacional da Serra do Divisor		12
Reserva Extrativista Cazumbá- Iracema	1,259	583
Reserva Extrativista Chico Mendes	42,795	8,607
Reserva Extrativista do Alto Juruá		206
Reserva Extrativista do Alto Tarauacá		95



Table S4. Areas of forest fires by municipalities in the state of Acre in the 1984 - 2016 period, classified by the number of times each pixel had been burned.

Municipality	Area burned (ha)		
	1 <sup>st</sup> time burned	2 <sup>nd</sup> time burned	3 <sup>rd</sup> time burned
Acrelândia	32,437	5,132	885
Assis Brasil	4,676	36	
Brasiléia	14,222	206	10
Bujari	24,058	4,886	70
Capixaba	12,838	2,600	305
Cruzeiro do Sul	916	11	2
Epitaciolândia	13,934	162	
Feijó	1,180	24	
Jordão	190		
Mâncio Lima	1,161	27	
Manoel Urbano	1,721	5	
Marechal Thaumaturgo	340		
Plácido de Castro	31,500	8,301	280
Porto Acre	23,526	3,410	201
Porto Walter	91		
Rio Branco	96,530	16,997	2,005
Rodrigues Alves	546	13	
Santa Rosa do Purus	2,402	9	
Sena Madureira	42,626	3,960	514
Senador Guimard	36,915	10,234	1,439
Tarauacá	270	5	
Xapuri	46,107	3,782	16

In Figure S1 one can see that August and September have large spikes of burning (active fire), most of which is in cattle pastures or agricultural fields but that trigger forest fires. This means that the forests are still burning in this period and it is not possible to map the total fire damage at that time. The best images for mapping the total fire damage in the forest are from September to December, when the fires decrease and the rains start.




Figure S1. Temporal distribution of hot spots between June and November in the period from 2000 to 2016 using the reference satellite Aqua-TM

The main challenge for mapping forest fires in the southwestern Amazon is confusion of fire scars with areas affected by blowdowns, vegetation under extreme water stress and senescent bamboo populations in open forests (Figure S2). These regional specificities can lead to errors in mapping land-use and land-cover change. In 2003, for example, INPE's PRODES project misclassified patches of senescent bamboo as deforestation (Deus and Mastrângelo, 2016), and, more recently, the MapBiomias project misclassified these same forest formations as secondary forests (MAPBIOMAS, 2016). The interpreter's local experience and knowledge is crucial to producing an accurate map.





Figure S2. Main types of confusion during the mapping of forest fires. (a) Bamboo senescence demarcated by black circles and (b) blowdowns near areas of fire marked by black rectangles and (c) vegetation under extreme water stress near large areas of forest fire.

1



2

3

4

Figure S3. Annual spatial distribution of the Maximum Cumulative Water Deficit (MCWD) anomaly for the State of Acre from 1998 to 2016.

6

7

8

9