

## Macro and microscopical identification of four species of *Baccharis* from trimera group

Budel, J.M.<sup>1</sup>; Duarte, M.R.<sup>1\*</sup>; Santos, C.A.M.<sup>1</sup>; Cunha, L.M.<sup>2</sup>

<sup>1</sup>Departamento de Farmácia, Universidade Federal do Paraná

<sup>2</sup>Curso de Farmácia, Universidade Federal do Paraná

### Abstract

Different species of *Baccharis* belonging to the Trimeria group are commonly named as *carqueja* and employed for their digestive and diuretic properties. Since the members of this group are alike, this work has aimed to study the macro and microscopical characters of *Baccharis articulata*, *B. cylindrica*, *B. gaudichaudiana* and *B. trimera*, in order to contribute for the drug analysis and to complement taxonomic knowledge for the Trimeria group. These species are shrubs and have three-winged stem, except for *B. articulata* which possesses two wings. *B. gaudichaudiana* axis reveals ribs, *B. articulata* has oval contour and *B. cylindrica* and *B. trimera* stem is circular. In *B. cylindrica* and *B. trimera*, included phloem has been observed in older basal regions. Calcium oxalate crystals are seen in the four species, being styloid in *B. articulata* and *B. cylindrica* and prismatic in *B. gaudichaudiana* and *B. trimera*. The other microscopical features observed are similar.

### Resumo

Diferentes espécies de *Baccharis* pertencentes ao grupo Trimeria são comumente denominadas como *carqueja* e empregadas por suas propriedades digestivas e diuréticas. Como os membros desse grupo são semelhantes, este trabalho objetivou estudar as características macro e microscópicas de *Baccharis articulata*, *B. cylindrica*, *B. gaudichaudiana* e *B. trimera*, como forma de contribuir à análise das drogas e complementar o conhecimento taxinômico do grupo Trimeria. Estas espécies são arbustos e tem caules tri-alados, com exceção de *B. articulata* a qual possui duas alas. O axis de *B. gaudichaudiana* revela nervuras, de *B. articulata* tem contorno oval e de *B. cylindrica* e *B. trimera* stem são circulares. Em *B. cylindrica* e *B. trimera*, floema incluso tem sido observado em regiões

basais antigas. Cristais de oxalate de calico são vistos nas quatro espécies, do tipo estilóide em *B. articulata* e *B. cylindrica* e prismáticos em *B. gaudichaudiana* e *B. trimera*. As outras características microscópicas observadas são similares.

The genus *Baccharis* comprises approximately 500 species, classified in 28 groups based on morphological similarities<sup>1</sup>. Different species belonging to the Trimeria group are commonly named as *carqueja* and employed for their digestive and diuretic properties<sup>2,3,4,5</sup>. In Brazil, the *carquejas* are one of the most commercialized medicinal plants and Paraná State is considered the major collector<sup>6</sup>. The Trimeria group members are alike and their identification presents some difficulties, even for specialists. Therefore, this work has aimed to study the macro (morphological) and microscopical (anatomical) characters of four species from this group, *Baccharis articulata* (Lam.) Persoon, *B. cylindrica* (Less.) DC., *B. gaudichaudiana* DC. and *B. trimera* (Less.) DC., in order to contribute for the drug analysis and to complement taxonomic knowledge for the Trimeria group.

The four species are shrubs, having winged stems which assume the leaf photosynthetic role, as foliar appendages are absent. In common, these species show some microscopical characters, such as uniseriate epidermis with polygonal shaped cells in face view and coated by thin and striated cuticle, anomocytic stomata even or slightly raised regarding the other epidermal cells, and clusters of pluricellular glandular trichomes united at the basis and localized in a small depression. The wings, in transection, reveal the chlorenchyma composed of an atypical palisade parenchyma beneath the epidermis and spongy parenchyma in the middle. Collateral vascular bundles are embedded in the chlorenchyma, encircled by a parenchymatic sheath and associated with secretory ducts. The caulinar axis has a dermal system similar to the wings, besides alternating chlorenchyma and angular collenchyma, and a vascular cylinder formed by phloem outside and xylem inside. The former may exhibit perivascular fibre caps and the latter surrounds the parenchymatic pith.

In contrast, *B. articulata* has a two-winged stem while the three others possess three wings. In transection, *B. gaudichaudiana* axis reveals ribs, *B. articulata* has oval contour and *B. cylindrica* and *B. trimera* stem is circular. In *B. cylindrica* and *B. trimera*, included phloem has been observed in older basal regions. Calcium oxalate crystals are seen in the four species, being styloid in *B.*

*articulata* and *B. cylindrica*, and prismatic in *B. gaudichaudiana* and *B. trimera*.

In conclusion, the macro and microscopical characters encountered for the four species correspond to the aspects described for the Trimeria group<sup>7,8,9,10</sup>. However, since outstanding characters lack for each species, the whole assembled characters are relevant for the pharmacognostic and taxonomic species identification.

### Material and Methods

The botanical material was collected in Paraná State and the vouchers were identified by a taxonomist. Stem fragments were fixed in FAA70<sup>11</sup>, maintained in 70% ethanol<sup>12</sup>, free-hand sectioned and stained either with toluidine blue<sup>13</sup> and basic fuchsin and astra blue<sup>14</sup>.

### Acknowledgments

The authors are thankful to Nelson I. Matzenbaker and Inês J. M. Takeda for the species identification.

### References

- <sup>1</sup>Barroso, G.M. Compositae - Subtribo Baccharidinae Hoffmann. Estudo das espécies ocorrentes no Brasil. *Rodriguésia*, v.28, p.1-273, 1976.
- <sup>2</sup>Corrêa, M.P. *Dicionário das plantas úteis do Brasil e das exóticas cultivadas*. Rio de Janeiro: IBDF, v.II, 1984; 707p.
- <sup>3</sup>Alonso, J.R. *Tratado de fitomedicina: bases clínicas y farmacológicas*. Buenos Aires: Isis, 1998; 1039p.
- <sup>4</sup>Takeda, I.J.M.; Farago, P.V. *Vegetação do parque estadual de Vila Velha: guia de campo*. Curitiba: Serzgraf, 2001; 419p.
- <sup>5</sup>Mors, W.B.; Rizzini, C.T.; Pereira, N.A. *Medicinal plants of Brazil*. Michigan: Reference Publications, 2000; 501p.
- <sup>6</sup>Correa Júnior, C; Ming, L.C.; Scheffer, M.C. *Cultivo de plantas medicinais, condimentares e aromáticas*. Curitiba: Emater-Paraná, 1991; 151p.
- <sup>7</sup>Spinar, L.A. Las especies de *Baccharis* (Compositae) de Argentina Central. *Boletim de La Academia Nacional de Ciências*, v.50, p.176-305, 1973.
- <sup>8</sup>Cortadi, A.; Di Sapia, O.; Mc Cargo, J.; Scandizzi, A.; Gattuso, S.; Gattuso, M. Anatomical studies of *Baccharis articulata*, *Baccharis crispa* e *Baccharis trimera*, "Carquejas" used in folk medicine. *Pharmaceutical Biology*, v.37, n.5, p.357-365, 1999.
- <sup>9</sup>Ortins, G.M.M.; Akisue, G. Estudo morfo-histológico, *screening* fitoquímico, constantes físicas e análise cromatográfica da droga e extrato fluido visando controle de qualidade da espécie *Baccharis articulata* Pers. *Lecta*, v.18, n.2, p.9-32, 2000.

<sup>10</sup>Sá, M.F.A.; Neves, L.J. Contribuição ao estudo das plantas medicinais *Baccharis myriocephala* DC. *Revista Brasileira de Farmácia*, v.77, n.3, p.88-96, 1996.

<sup>11</sup>Johansen, D.A. *Plant microtechnique*. New York: MacGraw Hill Book, 1940; 523p

<sup>12</sup>Berlyn, G.P.; Miksche, J.P. *Botanical microtechnique and cytochemistry*. Eames: Iowa State University, 1976; 326p

<sup>13</sup>O'Brien, T.P., Feder, N.; McCully, M.E. Polychromatic staining of plant cell walls by toluidine blue *O. Protoplasma*, v.59, n.2, p.368-373, 1965.

<sup>14</sup>Roeser, K.R. Die Nadel der Schwarzkiefer-Massenprodukt und Kunstwerk der Natur. *Mikrokosmos*, v.61, n.2, p.33-36, 1962.

### \* Autora para correspondência

Profa. Dra. Márcia do Rocio Duarte  
Laboratório de Farmacognosia - Departamento de Farmácia  
Universidade Federal do Paraná  
Rua Prefeito Lothário Meissner 3400  
Jardim Botânico  
CEP 80210-170 - Curitiba - PR  
Email: marciard@ufpr.br